

Holes and Shells in the Interstellar Medium of the Nearby Dwarf Galaxy IC 2574

Fabian Walter¹

Radioastronomisches Institut der Universität, Auf dem Hügel 71, D-53121 Bonn, Germany

Electronic mail: walter@astro.uni-bonn.de

and

Elias Brinks

Departamento de Astronomía, Apartado Postal 144, Guanajuato, Gto. 36000, Mexico

National Radio Astronomy Observatory, P.O. Box O, Socorro, NM 87801

Electronic mail: ebrinks@astro.ugto.mx

ABSTRACT

We present H I synthesis observation of the nearby dwarf galaxy IC 2574 (a member of the M81 group of galaxies) made with the NRAO Very Large Array in its B-, C- and D-array configurations at high spatial and velocity resolution ($95 \text{ pc} \times 2.6 \text{ km s}^{-1}$). In addition, we present optical broad and narrow band images obtained with the 2.2-m telescope of the Calar Alto observatory. The VLA H I observations show a stunning amount of detail in the form of 48 mostly expanding H I shells and holes in its neutral interstellar medium. These features range in size from about 100 (a limit set by the size of the beam) to about 1000 pc dominating the appearance of the H I surface brightness map. Their dynamics clearly influence the velocity field of IC 2574. In addition to well defined holes, some large scale ($> 1000 \text{ pc}$) coherent features are visible in the channel maps. They are probably the remainder of an older shell population.

¹Visiting Astronomer, German-Spanish Astronomical Centre, Calar Alto, operated by the Max-Planck-Institute for Astronomy, Heidelberg, jointly with the Spanish Commission for Astronomy.

Current star formation, as traced by H α emission, is predominantly found along the rims of the larger H I holes, suggesting propagating star formation. On linear scales of ≈ 95 pc, star formation occurs if the H I surface density reaches values higher than 10^{21} cm $^{-2}$. The radial expansion of the H I holes (8–12 km s $^{-1}$), the indicative ages (10–60 10^6 yr) and the energy requirements for their formation (10^{50} – 10^{53} ergs) can be understood in terms of the combined effects of stellar winds and multiple supernova explosions of the most massive stars formed during a recent phase of active star formation.

The scaleheight of the H I layer is found to be ≈ 350 pc, considerably thicker than that in massive disk galaxies. This is due to a lower gravitational potential (for the same observed onedimensional velocity dispersion of ≈ 7 km s $^{-1}$). This puffed up disk implies a lower H I volume density (0.15 cm $^{-3}$). This, combined with the reduced gravitational pull and solid body rotation throughout the disk explains why the diameter distribution in dwarf galaxies extends to substantially larger values than in spiral galaxies.

A comparison with other galaxies shows that the energies needed to create these structures are the same for all types of galaxies, at least to first order. The overall statistical properties of the H I holes and shells in galaxies show clear trends with Hubble type (or rather mass), such as in their diameter distribution, expansion velocities and ages.

Subject headings: Galaxies: individual (IC 2574) – Galaxies: ISM – ISM: structure – Radio lines: galaxies – Radio lines: ISM

1. Introduction

Some twenty years ago, Heiles (1979, 1984, 1990) discovered large, shell-like structures in the neutral hydrogen (H I) distribution of our Galaxy by inspection of maps made in the 21-cm line. Since this discovery, a wealth of observations obtained with powerful Synthesis Radio Telescopes such as the Very Large Array (VLA) or the Westerbork Synthesis Radio Telescope (WSRT) revealed similar structures in our nearest neighbors. Prominent examples are M 31 (Brinks 1981, Brinks & Bajaja 1986), M 33 (Deul & den Hartog 1990), Holmberg II (Puche *et al.* 1992, hereafter referred to as PWBR92), the galaxies M 101 and NGC 6946 (Kamphuis 1993) and IC 10 (Wilcots & Miller 1998). The latest, impressive additions to this list are the SMC (Staveley-Smith *et al.* 1997) and LMC (Kim *et al.* 1998). All these observations indicate that the interstellar medium (ISM) of medium to late-type galaxies is dominated by features, which are variously described as shells, rings, holes, loops, bubbles or cavities.

Obviously, one of the basic questions is which physical process is able to produce these observed structures. The general picture which has emerged is that they are the result of combined stellar winds and supernova explosions produced by young stellar associations. For review articles on that topic see Tenorio-Tagle & Bodenheimer (1988), and van der Hulst (1996, and references therein). Based on a simple model of holes created by O and B stars, Oey & Clarke (1997) successfully predict the observed number distribution of SMC holes, lending support to this hypothesis. It should be noted, however, that this general picture is not without its flaws. Searches for the remnant stellar populations of the OB associations thought to be respon-

sible for some of the holes in Ho II have not led to the expected result (Rhode *et al.* 1999). And in the case of the largest observed shells, these seem to have energy requirements surpassing the output of stellar winds and supernovae. To explain those structures an alternative mechanism was proposed, the infall of gas clouds (see Tenorio-Tagle *et al.* (1987) for a numerical simulation and van der Hulst & Sancisi (1988) for observational evidence in the case of one of the largest holes in M 101). And recently, even Gamma Ray Bursters (GRBs) have been suggested as a possible explanation for the largest H I holes (Efremov, Elmegreen & Hodge 1998, Loeb & Perna 1998).

The idea that supernovae deposit large amounts of energy into the ISM is not new. Cox & Smith (1974) already discussed the idea that the energy released by supernovae sets up tunnels of hot gas in the ISM. This picture was a substantial modification of the Field, Goldsmith & Habing (1969) model which consists of a two phase medium. The ideas proposed by Cox & Smith were incorporated within the model elaborated by McKee & Ostriker (1977) who suggested that the ISM has three distinct phases, a cool ($T \approx 10^2$ K), a warm ($T \approx 10^4$ K), and a hot phase ($T \approx 10^6$ K), all in pressure equilibrium. Various refinements have been proposed but this picture basically still stands.

Knowing how much kinetic energy is transferred in expanding shells and filaments is of direct importance for assessing the energy balance in the ISM. In addition to an analytical approach, various authors are pushing forward numerical simulations (e.g., Palouš, Franco & Tenorio-Tagle 1990, Silich *et al.* 1996, Palouš, Tenorio-Tagle & Franco 1994), in order to model the evolution of the shape

of the H I shells in various environments, the fragmentation of matter on the rim of the shells and to investigate induced star formation. This is important not only for tracking down the mechanisms which produce the H I shells and filaments, but also for determining what fraction of the ISM breaks out into the halo and might eventually rain back onto the disk.

Our aim is to investigate the properties of the H I shells and bubbles in the ISM of galaxies. Intimately linked to this are the questions when fragmentation on the rim of the shells starts and what drives star formation. Unfortunately, the determination of the characteristics of H I shells in our Galaxy is not straightforward due to our unfavorable location within the disk. Looking at the nearest spiral galaxies (e.g., M 31, M 33, M 101, NGC 6946) does not improve matters all that much since the structures in their ISM are strongly modified by spiral density waves and differential rotation, both effects dramatically influencing the shape and formation of holes and shells. Therefore, we decided to direct our attention to nearby dwarf galaxies instead. Dwarfs are slow rotators, generally show solid body rotation, and lack density waves. This implies that once features like shells have formed, they won't be deformed by galactic shear and therefore tend to be long lived. Moreover, the overall gravitational potential of a dwarf is much smaller than in a normal spiral. The same amount of energy input of a star forming region therefore has a more pronounced impact on the overall appearance of the ISM, as shown by PWBR92. Since dwarf galaxies also tend to have a much thicker disk compared to massive spirals (e.g., Staveley-Smith, Davies & Kinman 1992), the H I volume density of dwarf galaxies is lower,

again facilitating the creation of large holes. And because of their smaller size, the probability to suffer impacts from infalling clouds is lower compared to that for larger galaxies. Despite all these arguments in favor, only very few such H I studies of dwarf galaxies have been published. Beside our two neighbors, the SMC and LMC a detailed analysis was performed in one case only, that of the dwarf galaxy Holmberg II (PWBR92), member of the M 81 group.

In this paper, we concentrate on IC 2574, a gas-rich dwarf galaxy in the same group. Table 1 summarizes some general information on this object. IC 2574 (= UGC 5666 = DDO 81, also sometimes referred to as Coddington's Nebula) has been studied by several authors in the past. Observations with single dish telescopes in the 21-cm line of neutral hydrogen (H I) were published by Rots (1980) and Huchtmeier & Richter (1988) and a first study using an interferometer was made by Seielstad & Wright (1973). A recent H I study, concentrating on the dynamics and contribution to the total mass by dark matter was performed by Martimbeau, Carignan & Roy (1994, hereafter abbreviated MCR94), who observed the galaxy with the Westerbork Synthesis Radio Telescope. Recent broadband optical studies were published by Tikhonov *et al.* (1991). Studies of H II regions in IC 2574 have been presented by Miller & Hodge (1994), Kennicutt (1988) and Arsenault & Roy (1986). Spectroscopy of some prominent H II regions was performed by Miller & Hodge (1996) and H α velocity fields were recently obtained by Tomita *et al.* (1998). In addition, four Wolf-Rayet stars were discovered by Drissen, Roy & Moffat (1993).

We present H I observations made with the

NRAO² Very Large Array (VLA) in its B-, C- & D-array configurations supplemented by optical observations obtained at the Calar Alto Observatory. The VLA data were originally obtained by Puche, Brinks and Westpfahl with the intention of following up on their work on Ho II. So far, only a low resolution integrated H I map was published by Puche & Westpfahl (1994). This paper presents the VLA data in a complete form and compares them with optical observations. The original data were recalibrated by us to produce far superior maps using the latest calibration and imaging techniques within AIPS. The data reduction and results are discussed in Secs. 2 & 3. In Sec. 4, a catalog of H I holes and shells is presented. Sec. 5 discusses the relation between H I and H α features. A general discussion is given in Sec. 6 and a summary of the results is presented in Sec. 7.

2. Observations

2.1. Radio Observations

IC 2574 was observed with the VLA in B-, C- and D-configuration. Part of the D-array observations were affected by solar interference, which was removed by deleting uv -spacings shorter than typically 0.4 k λ and editing out bad time-ranges. In total, 17 hours were spent on source which were divided into 4 hours for D-array, 2.5 hours in C-array and 10.5 hours in B-array (see Table 2 for a detailed description of the VLA observations). The longer observing time in B-array only partly compensates for the lower surface brightness sensitivity in

this extended configuration. The absolute flux calibration was determined by observing 1328+307 (3C286) for approximately 20 minutes during each observing run, assuming a flux density of 14.73 Jy according to the Baars *et al.* (1977) scale. This calibrator was also used to derive the complex bandpass corrections. The nearby calibrators 1031+567 and 0945+664 were used as secondary amplitude and phase calibrators and their fluxes were determined to be 1.80 Jy and 2.22 Jy, respectively. Since the systemic velocity of IC 2574 (≈ 53 km s⁻¹) is close in velocity to H I emission from our Galaxy, each calibrator observation was split in two parts to form a pair of observations, one part having the velocity shifted by +300 km s⁻¹ and the other by -300 km s⁻¹ to avoid contamination by Galactic emission. In the course of the calibration these observations were then averaged to give interpolated amplitude and phase corrections. IC 2574 was observed using a 1.56 MHz bandwidth centered at a heliocentric velocity of 38 km s⁻¹. This band was divided into 128 channels resulting in a velocity resolution of 2.58 km s⁻¹ after online Hanning smoothing.

The data for each array were edited and calibrated separately with the AIPS package³. The uv -data were inspected and bad data points due to either interference or cross talk between antennae were removed, after which the data were calibrated. We Fourier transformed our B-, C- and D-array observations separately to assess their quality; subsequently we combined all data to form a single dataset which was used for mapping. All the results presented in this paper are based on this combined dataset.

Channels with velocities lower than -40

²The National Radio Astronomy Observatory (NRAO) is operated by Associated Universities, Inc., under cooperative agreement with the National Science Foundation.

³The Astronomical Image Processing System (AIPS) has been developed by the NRAO.

km s^{-1} and higher than $+136 \text{ km s}^{-1}$ were found to be line free; these channels determined the continuum emission which was subtracted in the uv -plane. The same channels were used to produce a radio continuum map of IC 2574.

After that, two sets of datacubes (1024×1024 pixels $\times$ 68 channels each) were produced using the task IMAGR in AIPS, each of them CLEANed to a level of two times the rms noise (Högbom 1974, Clark 1980). The first data set was made with natural weighting, leading to a resolution of $12.4'' \times 11.9''$ and emphasizing large scale structures. A second cube was produced using the ROBUST weighting scheme (Briggs 1995). This scheme achieves the high sensitivity of natural weighting combined with a well behaved synthesized beam, at a resolution close to that of uniform weighting. To obtain the optimum ROBUST value for our observation, the ROBUST parameter space was searched by producing maps in the sensitive regime of $-2 < \text{ROBUST} < 2$ and measuring for each map the rms noise and beamsize. Eventually, we chose a value of $\text{ROBUST}=0.25$, resulting in a beamsize of $6.4'' \times 5.9''$ and an rms noise of $0.7 \text{ mJy beam}^{-1}$. A comparison with the beamsize of the high resolution uniform map ($4.7'' \times 4.3''$) and the rms noise of the low-noise, natural map ($\approx 10\%$ increase in noise) shows the strength of this mapping algorithm. Unless otherwise mentioned, the results presented in the following are based on this ROBUST cube.

Although IC 2574 is situated at a galactic latitude of 43.6° , some of the channels were clearly confused by Galactic emission (in particular between $+10$ and -10 km s^{-1}). Since galactic emission shows its presence on predominantly large scales, we were able to re-

move the most prominent emission features by blanking the uv -data out to $0.1 \text{ k}\lambda$ in these channels.

To separate real emission from the noise, the following procedure was applied: the natural weighted data cube was convolved to a circular beam with a FWHM of $45''$. The smoothed map was then tested at the 2σ level; if a pixel fell below this level, the counterpart in the cube was blanked. After that, the remaining peaks were inspected. Emission that was present in 3 consecutive channels was considered to be real while all other remaining spikes were considered to be noise and blanked. The final result was named the master cube. This master cube can be used as a mask to blank the natural and ROBUST data cubes. This method ensures that the same regions are included when inspecting cubes at different resolutions and with different signal-to-noise ratios.

2.2. Optical Observations

IC 2574 was observed with the 2.2-m telescope of the Calar Alto Observatory⁴ on 1999 January 25 and 26 in Johnson R-band (20 min) as well as in $\text{H}\alpha$ (45 min). A 2048×2048 pixel CCD was used leading to $0.5''$ per pixel. The seeing was around $1.3''$. A focal reducer (CAFOS) allowed for a field of view of radius $8'$ to be imaged. The usual calibration steps were followed using the IRAF package⁵. Each exposure was corrected for zero offset (bias) and was flatfielded using skyflats. Af-

⁴The Calar Alto Observatory is operated by the Max-Planck-Institute for Astronomy, Heidelberg, jointly with the Spanish Commission for Astronomy.

⁵IRAF is distributed by National Optical Astronomy Observatories, which is operated by the Association of Universities for Research in Astronomy (AURA), Inc., under contract with the National Science Foundation.

ter that, frames were inspected for bad pixels and shifted to get them aligned. Lastly, the exposures were combined to eliminate cosmic ray hits and to create a final image. As it turned out, conditions were not photometric during the observation, hence no calibration using standard stars was performed. In order to get an approximate calibration for IC 2574, values were taken from the literature (MCR94). The deep R-band image was used for the continuum subtraction of the H α map. The continuum level was determined by simply scaling the continuum image relative to the line plus continuum image such that the foreground stars disappeared when the two images were subtracted. For the calibration we relied on the H α fluxes published by Miller & Hodge (1994, hereafter referred to as MH94). In their paper, these authors identify 289 H II regions in IC 2574 (for a thorough discussion on the H α emission and its calibration see the Appendix in MH94). Figs. 1 and 2 show our R-band and the H α image, respectively.

Coordinates for the maps were obtained using the task KOORDS within the KARMA software package (for a description of KARMA see Sec. 4.1). We used as input an image from the Digital Sky Survey. Using 14 stars found in our CCD image, it was possible to determine the coordinate system to an accuracy of better than one arcsecond, more than adequate for our needs.

3. HI Studies

3.1. Global H I Properties

The channel maps obtained by the robust weighted cube are presented in Fig. 3. As can be seen, most traces of contamination by Galactic emission were successfully removed.

To save space, only every other channel over the frequency range where line emission is present is shown. The heliocentric radial velocities (in km s^{-1}) are plotted in the upper right-hand corner of each plane. The beam-size ($6.4'' \times 5.9''$) is indicated in the top left-hand panel (note that the beamsize is so small that it cannot be properly reproduced in our maps). Many holes and shell-like structures are visible in the channel maps; these features will be the main topic of this paper.

Obtaining the H I flux of each channel and hence the global H I profile of IC 2574 is not straightforward. As first pointed out by Jörsäter & van Moorsel (1995), fully cleaned maps do not exist and any cleaned map consists of the sum of two maps: one containing the restored clean components and the other the residual map. In the former, the unit is Jansky per clean beam area, and in the latter Jansky per dirty beam area. Usually, fluxes are determined on a cleaned map, assuming that the clean beam is the correct one for the entire map. In reality, the flux is calculated correctly only for the cleaned fraction of the map; for the residual map the flux is overestimated when the dirty beam area is bigger than the clean beam area (which is usually the case). This becomes a real problem for extended objects, especially in the case of combined array data, like for our VLA data set. For a full discussion on this topic see Jörsäter & van Moorsel (1995). Following their prescription, the real flux of a channel map is given by:

$$G = \frac{D \times C}{D - R}$$

where C is the cleaned flux, and R and D are the 'erroneous' residual flux and the 'erroneous' dirty flux over the same area of a channel map, respectively. We followed this

procedure for every single channel to obtain the global H I profile as presented in Fig. 4. Without this correction, we would have overestimated the H I flux by a factor of about two!

Figure 4 shows the flux- and primary-beam corrected fluxes for every channel of IC 2574 (open squares) plotted together with the single dish profile published by Rots (1980) and the WSRT data by MCR94 (the arrows indicate the velocity range for which the data points were interpolated due to contamination by galactic emission). From our data, we derive a total integrated H I flux of 373 Jy km s^{-1} which should be compared with the value of 399 Jy km s^{-1} obtained by Rots (1980), who used the Greenbank 300 ft telescope to map IC 2574. This is excellent agreement, given the uncertainties of removing Galactic emission and systematic errors in the absolute calibration. This also means that we hardly miss any H I flux due to missing short spacings. Other values obtained by interferometric observations are: $440 \pm 50 \text{ Jy km s}^{-1}$ found by Seielstad & Wright (1973), using the Owens Valley Radio Interferometer and 286 Jy km s^{-1} obtained by MCR94 from their Westerbork observations. As can be seen from Fig. 4, the new VLA observations recover substantially more flux than the WSRT data. Adopting a distance of 3.2 Mpc, our flux translates to a total neutral hydrogen mass of $M_{HI} = (9.0 \pm 0.1) \times 10^8 M_{\odot}$ for IC 2574.

3.2. Global Characteristics

Fig. 5 shows the H I surface density map of IC 2574 obtained by adding all channels containing H I emission of the robust cube, after blanking them with the master cube. Note that the beamsize is so small that it

can hardly be properly reproduced. Note also that, although many hole-like structures and cavities are visible, specific features turn out to be less prominent in the total surface brightness map than in the channel maps. This is related to the fact that the galaxy is highly inclined, leading to overlapping of shell-like structures along the line-of-sight in the integrated map. For a more complete discussion on that topic, see Sec. 4.1.

Fig. 6 shows the velocity field of IC 2574 which was calculated in the usual manner by taking the first moment. Since the high resolution velocity field is dominated by small-scale motions within the ISM, the natural weighted cube was convolved to $30''$ (as indicated by the beamsize) to put more emphasis on the global characteristics of the velocity field. The underlying greyscale is a linear representation of the H I surface brightness map at full resolution. Although the overall velocity field shows the typical behavior for a dwarf galaxy (solid body rotation in the center, flattening of the rotation curve towards the very edge of the galaxy), the general rotation is obviously disturbed by noncircular motions (e.g., note the prominent band of lower velocities to the north east of the center of the galaxy, parallel to its major axis). These deviations are linked to holes in the neutral interstellar medium.

Finally, Fig. 7 shows the second moment map, or velocity dispersion of IC 2574. Although some regions near shells reach values as high as 15 km s^{-1} , the overall velocity dispersion in the unperturbed parts is much lower. Using H I profiles, we derive the velocity dispersion in the disk by looking at quiescent regions. Several lines of sight were averaged to give a one sigma velocity dispersion of $7 \pm 1 \text{ km s}^{-1}$ (similar to what is found in

Ho II). This shows that our velocity resolution of 2.5 km s^{-1} (FWHM) was quite sufficient to resolve the lines of neutral hydrogen.

In this paper, we focus on the small-scale structure of the ISM in IC 2574. For our analysis, however, we need to adopt values for the orientation of the object which are usually derived on the basis of a kinematical analysis (rotation curve fitting). A full analysis of the dynamics of IC 2574 based on our observations, however, is beyond the scope of this paper. Luckily, an adequate analysis has already been published (see the paper by MCR94). In the course of their study they derive a systemic velocity of $58 \pm 7 \text{ km s}^{-1}$, adopt an inclination of $75^\circ \pm 7^\circ$ and a position angle of 52° . Depending on their dynamical model, they find that 70 – 90% of the total mass of the galaxy is in the form of dark matter. Following the same procedure as performed for the dwarf galaxy II Zw 33 (Walter *et al.* 1997) we carried through a first analysis of our data confirming the results by MCR94 regarding the rotation curve and the global orientation of IC 2574. For ease of comparison we therefore adopted their values throughout this paper.

3.3. The scale height of the H I disk

The scale height h of the H I layer of IC 2574 plays an important role in the determination of the properties of the holes, e.g., when translating the observed H I surface density into an H I volume density. Here h is defined as the 1σ value for a Gaussian distribution of the H I layer. Recent studies have shown that in general dwarf galaxies have thicker H I disks than more massive spiral galaxies. Their aspect ratio tends to be of the order of 10:1 rather than 100:1, e.g., Ho II: $h = 625 \text{ pc}$ (PWBR92) or NGC 5023:

$h = 460 \text{ pc}$ (Bottema *et al.* 1986; see also Staveley-Smith *et al.* 1992). In the case of IC 2574 there are also indications that the H I is distributed in a thicker disk. MCR94 already noticed that, looking at the apparent axial ratio of the integrated H I map and assuming a real inclination of 75° , the H I must be distributed in a somewhat thicker disk. Moreover, the fact that we observe, as we will argue below, spherically expanding shells with diameters of up to almost 1 kpc implies that the extent of the H I layer must be of the same order of magnitude. Hence, the scale height of the H I gas should be of order 350–400 pc.

An approximate value for the scale height can be obtained fairly simply, as shown by PWBR92. We follow their approach here. The scale height is proportional to the velocity dispersion of the gas and inversely proportional to the square root of the mass density in the disk (Kellmann 1972; van der Kruit 1981). For a density distribution that can be described as $\rho(z, R) = \rho(0, R) \text{sech}^2(z/z_0)$, z_0 is given by

$$z_0(R) = \frac{\sigma_{gas}}{\sqrt{2\pi G \rho(0, R)}}$$

where R is the galactocentric distance, σ_{gas} the velocity dispersion of the gas in quiescent regions of the galaxy (Sec. 3.2), and $\rho(R)$ the total volume density (gas, stars and dark matter) of the galaxy in the plane. However, since $\text{sech}^2(z/z_0) \approx \exp(-z^2/z_0^2)$ (to within 5% in the region of interest) we can also assume a Gaussian distribution in z . The 1σ scale height h is then defined as $h = z_0/\sqrt{2}$. To calculate the average density of IC 2574 in the plane we assume that the total mass of the galaxy can be estimated from the last measured point (R_{max}) of the observed ro-

tation curve (MCR94) and that this is distributed over a spherical volume of radius R_{max} . MCR94 derive a rotation curve with $v(R_{max}) = 66 \text{ km s}^{-1}$ at $R_{max} = 8.0 \text{ kpc}$. This corresponds to a total dynamical mass of $8.3 \times 10^9 M_{\odot}$. Distributing this mass over a spherical volume with radius R_{max} gives an average value for ρ of $0.0039 M_{\odot} \text{ pc}^{-3}$. Since this is only the mean density of the galaxy and the density near the plane is certainly higher, we take twice this derived value (see PWBR92 for a justification) to get a first approximation to the real volume density $\rho(R)$ near the center. Using this approach, we find $\rho(R) = 0.008 M_{\odot} \text{ pc}^{-3}$.

Substituting the observed velocity dispersion of 7 km s^{-1} (see Sec. 3.2) in the equation given above we derive a scale height of $h = 345 \text{ pc}$ for IC 2574. Note that this value, given all the uncertainties in the derivation, is in good agreement with the value derived based on the sizes of the largest holes ($h = 350 - 400 \text{ pc}$). We therefore adopted a scale height of $h = 350 \text{ pc}$ for the H I layer of IC 2574.

We can now calculate the volume density of the H I from the projected and corrected to face-on H I surface density. The surface density profile was obtained by elliptically averaging the flux- and primary-beam corrected total H I map (assuming an inclination of 75°). Again, this is correct to first order only as in the case of a thick H I distribution the deprojection of the observed projected H I distribution is not trivial. We then calculated the volume density $n_{HI}(R)$ of the H I in the plane of the disk using

$$N_{HI}(R) = \int_{-\infty}^{+\infty} n_{HI}(R) \exp\left(\frac{-z^2}{2h^2}\right) dz = \sqrt{2\pi} h n_{HI}(R)$$

4. H I Holes in IC 2574

4.1. The H I-hole catalogue

We conducted a search for H I structures in IC 2574 using the tasks KVIEW, KPVS-LICE and KSHELL that are built into the recently developed KARMA⁶ visualization software package. KVIEW offers many different ways to look at a movie of a datacube whereas KPVS-LICE produces real-time position velocity (pV) cuts through a cube in any orientation which tremendously facilitates the search for and identification of hole-like structures. Finally, KSHELL was used to make radius-velocity plots of some of the more prominent shells and holes.

First, the ROBUST cube was searched for the smaller holes in IC 2574. In order to be sensitive to more extended structures and therefore bigger holes, we smoothed our ROBUST cube to $11''$, $15''$ and $30''$, respectively, and inspected each of them. For each H I hole in IC 2574 we determined its position, size and expansion velocity. Both authors independently produced a list of holes to reduce personal bias, based on some simple criteria regarding the determination of the center, size, expansion velocity and systemic velocity of the holes. After that, the two lists were compared and merged. There was an overlap of about 75% in terms of well defined holes which lends some credibility to the identification process. We reexamined the remaining structures and, erring on the side of caution, ended up with a catalogue of 48 holes. Although this list cannot be free from personal bias, the method closely follows that employed in the case of M 31 and Ho II which should make a comparison between those ob-

⁶The KARMA visualization software package was developed by Richard Gooch of the Australia Telescope National Facility (ATNF)

jects and IC 2574 valid.

It is interesting, at this point, to reflect on the reality of the features. Do they really constitute large-scale, coherent structures or are we tricked by nature into seeing coherence in what is basically a turbulent, perhaps fractal medium? Mac Low *et al.* (1998) recently produced sets of model data cubes, similar to those obtained via radio (H I) or optical (H α) observations (i.e., position, position and velocity). We analysed, using the same techniques as explained above, a cube of uniform, isotropic, isothermal, supersonic, super-Alfvénic, decaying turbulence. The simulations assume that the gas is optically thin and emits with a strength directly proportional to its density. The model corresponds to a weakly magnetized ISM. Single cuts through the data cube show features remarkably similar to the ones seen in the H I data. However, when focussing on a potential feature and investigating its morphology in position–position or position–velocity space, no coherent structure emerges. This lends us to believe that shells due to random (or at least turbulent) processes make up a small fraction of the holes we classified. Instead, it is more likely that the H I holes and shells are due to mechanisms which act on large scales, such as stellar winds and (multiple) supernovae.

Following Brinks & Bajaja (1986), we distinguished between three different types of holes based on their appearance in the pV–diagrams. A Type 1 hole corresponds to a total blowout (i.e., we don’t observe the receding nor the approaching side of the shell). In that case, it is impossible to determine the expansion velocity and hence the kinematics of the hole. Figure 8c shows a pV–diagram of a typical Type 1 hole (our hole number 30)

using KPVSLICE.

A Type 2 hole corresponds to a hole which is offset with respect to the plane of the galaxy. In that case, we witness a deformation in the pV–diagram towards the plane of the galaxy and a blowout to the opposite side (see Silich *et al.* 1996 for numerical simulations of such a case — their case C). The amount of deformation yields the expansion velocity of the shell towards the midplane. The part of the shell that points away from the plane is usually not visible due to lack of sensitivity. Figure 8d shows a pV–diagram of a typical Type 2 hole (our hole number 33).

Type 3 is the ‘classic hole’ and corresponds to a more or less complete expanding shell. The signature in pV–space is that of an elliptical hole. The expansion velocity is equal to half the difference between the velocity of the approaching and receding sides. The systemic velocity of the hole is defined by its center. Figure 8a shows a pV–diagram of a typical example of a Type 3 hole (our hole number 21). Figure 8b shows the same hole, but analyzed with the task KSHELL. In this figure, the horizontal axis corresponds to radius r (distance from the center of the hole) and the intensity corresponds to the average H I density within an annulus with radius r around the center of the hole. The vertical axis is again velocity. Since we are looking at a radius–velocity diagram, we will call such a representation an rV–diagram in the following. For an ideal shell, a half ellipse should be visible in which the diameter in the v –direction is twice the expansion velocity and its dimension in the r –direction the radius of the shell. The advantage of looking at a near circular H I shell with KSHELL lies in a substantial increase in signal-to-noise ratio as one averages over many pixels whereas in a pV–diagram

one merely cuts through an H I structure.

Note that in the case of a Type 1 hole, the absence of detectable emission corresponding to the front and back of a shell causes a hole in rV-space. For obvious reasons, a Type 2 hole does not show a prominent signature in an rV-diagram either.

As was already mentioned earlier, the holes were found to be much more prominent in the channel maps than in the integrated H I map. This is due to the high inclination of IC 2574 which leads to several H I shells at different locations within the disk and possibly different velocities overlapping along the line-of-sight, as is the case in M 31. The total H I surface brightness map was therefore of little use for the identification of a hole. Note that in the case of Ho II, *only* the integrated map was used because of the much lower inclination of the galaxy. This implies that the approach to search for holes needs to be adapted to the orientation of the galaxy.

Figure 9 shows a plot of the distribution of the 48 H I holes detected with confidence in IC 2574. The greyscale map is a linear representation of the total H I surface brightness. Some of the holes are clearly visible in the total H I map but the majority of holes doesn't show up at all and can only be detected in the channel maps or pV-diagrams.

The final list of the detected H I holes is given in Table 3. The properties listed are as follows:

- Column 1:** Number of the hole (increasing with increasing right ascension)
- Column 2, 3:** Position of the center of the hole in right ascension and declination, respectively (B1950.0 coordinates). Typical errors are 3".

Column 4: Heliocentric velocity v_{hel} of the hole defined by the channel in which the hole is largest or most prominent. The accuracy of v_{hel} is about 2.5 km s^{-1} (the velocity resolution of our observations).

Column 5: Diameter d of the hole in kpc. Note that most of the holes are circular. In the case of elliptical holes, the diameter along the major axis, along the minor axis and the position angle ($2 \times b_{\text{maj}}, 2 \times b_{\text{min}}, PA$) is given (but see Sec. 4.2). The accuracy of d is around 95 pc (the FWHM of our synthesized beam).

Column 6: The expansion velocity v_{exp} of the hole in km s^{-1} . Note that in the case of a Type 1 hole, no expansion velocity can be measured. Typical errors are about 2 km s^{-1} .

Column 7: The estimated H I volume density n_{HI} (in $\text{M}_{\odot} \text{ pc}^{-3}$) of the region where the hole is situated before the hole was created. n_{HI} was calculated based on the H I surface density, as described in Sec. 3.3. Given the uncertainties in the deprojection and the assumptions which went into the derivation of the mass density within the disk, we estimate a relative error in the value of n_{HI} to be of order 30%.

Column 8: The type of the hole (1–3), as defined in Sec. 4.1.

Table 4 summarizes the physical properties derived from Table 3. The properties listed are as follows:

Column 1: Number of the hole

Column 2: Diameter d of the hole. For elliptical holes, the diameter was defined using the geometric mean

$$d = \sqrt{2 b_{\text{maj}} \times 2 b_{\text{min}}}$$

Column 3: Expansion velocity v_{exp} of the hole in km s^{-1} . Same as column 6 in Table 3.

Column 4: Kinematic age of the hole in units of 10^6 years. This age determination assumes that over its entire lifetime the hole was expanding with v_{exp} . Hence:

$$t = \frac{(d/2)}{v_{\text{exp}}}$$

The accuracy of the kinetic age is typically 20%. Because of the assumption of a constant expansion velocity, the kinematic age is an upper limit.

Column 5: Indicative H I mass M_{HI} in units of $10^4 M_{\odot}$. It is the mass which filled the observed hole under the following assumptions: i) the hole is spherical, ii) the gas was distributed uniformly in the galaxy, and iii) the hole is completely empty now. Note that if one can assume that only a minor fraction of the surrounding H I is ionized, this mass corresponds also to the total H I mass on the rim of the hole. M_{HI} was calculated using

$$M_{\text{HI}} = \frac{4}{3} \pi (d/2)^3 n_{\text{HI}}(R)$$

where $n(R)$ is the volume density of the gas at the galactocentric distance R (column 7 in Table 3). To correct this value for the amount of primordial helium, a factor of 1.33 has to be applied.

The relative error is, as d enters to the third power, of order 30%. Note that in the case of the largest holes, this value is more likely to be an upper limit since n_{HI} decreases with increasing distance z from the plane.

Column 6: The initial total energy E_{E} that was released by the supernova explosions which created the hole (in units of 10^{50} ergs). Note that the kinetic energy of the expanding shell is about 1% to 10% of E_{E} (depending on the ambient density, see Chevalier 1974). To determine E_{E} , we used the equation by Chevalier (1974) who derived a fit to numerical results of an expanding shell created by a single explosion (see also McCray & Snow 1979). Expressing E_{E} in terms of observable quantities d , v_{exp} and n_0 , he obtained

$$E_{\text{E}} = 5.3 \times 10^{43} n_0^{1.12} (d/2)^{3.12} v_{\text{exp}}^{1.4} \text{ ergs}$$

where v_{exp} is the expansion velocity in km s^{-1} , n_0 the density of the ambient medium in particles per cubic centimeter, and d is the diameter of the shell in parsecs. In our calculations we replaced n_0 by n_{HI} as was done, e.g., for M 31 and Ho II. This is not strictly correct as we are ignoring contributions by He and heavier elements. To correct E_{E} for this contribution, all values should be multiplied by a factor of ≈ 1.5 . In addition, the derived H I densities are average values for a given galactocentric distance which introduces another source of uncertainty. Note that the numerical coefficient in Chevaliers equation depends on the radiative energy loss rate and thus on the heavy element abundance in the expanding shell. Not only

is this abundance different for various galaxies, but it decreases with galactic radius, both for our own and external galaxies and by more than an order of magnitude (e.g., Heiles 1979). The last phase in the expansion of a shell is reached when, after the shell has continued to collect more of the ambient medium it will slow down to velocities comparable to the random motions of the ISM (around 7 km s^{-1}) and hence stops expanding. Most probably the largest structures we observe are in this latter phase (see Sec. 4.3). From this brief discussion it should be clear that the derived energies E_E are only order of magnitude estimates.

4.2. Notes on single holes

The following holes deserve a more detailed description:

Hole #35: This hole causes a major disturbance in the pV-diagrams of IC 2574. It is one of the few elliptical holes, with dimensions of $1000 \times 500 \text{ pc}$. Although there is some circumstantial evidence that it is a superposition of two spherical holes, it was treated as one hole. As we will show in Sec. 5, the most prominent H II region in IC 2574 is located on the rim of this supergiant shell. The shell is filled with soft X-ray emission as observed with the ROSAT telescope. A detailed multiwavelength analysis (optical, radio, X-ray) of this particular supergiant shell has been presented elsewhere (Walter *et al.* 1998)

Hole #22: Catalogued elliptical in shape, this hole is actually a superposition of several holes, leading to the overall impres-

sion of an elliptical feature; we treated this as a single object since subdividing it in, say, three holes would have been arbitrary.

Holes # 13, 23, 26: As can be seen from the channel maps (Fig. 3), some holes are not clear-cut cases in terms of the definition of a hole given in Sec. 4.1. They are possibly chance superposition of old shells and fragments leading to the impression of a hole in the integrated H I map (see also Sec. 4.3). We therefore decided not to include these holes in the statistical analysis.

4.3. Notes on large-scale structures

As can be seen from Table 3, only few shells larger than $\approx 1000 \text{ pc}$ have been catalogued in IC 2574. It has to be mentioned, however, that there appear coherent features larger than 1 kpc in the channel maps. Several of these can be identified in the channel maps (Fig. 3), e.g., between 45 and 70 km s^{-1} . Note that these structures are related to the major disturbance in the velocity field due north of the center of the galaxy. Obviously, it is very difficult to catalogue these features as they lack the clear coherence, both in position as well as in velocity, of H I shells. Moreover, because of their large size, their dynamics is not dominated by some leftover expansion but rather by large scale galactic rotation modified by the effects of nearby, smaller holes. The reason why we wish to mention these large scale features is because along some of the apparent arcs we find H II regions.

5. The H I–H α connection in IC 2574

5.1. Global correlations

Until now, we have focussed on the structure and dynamics of the neutral ISM of IC 2574. In the following, we will compare these results with our H α observations which reveal the sites of recent or ongoing SF in IC 2574. As noted earlier (Sec. 1), the expanding H I structures are widely thought to result from recent star formation. They sweep up gas causing the volume density on the rim of the shell to increase until gravitational instabilities lead to molecular cloud formation and hence secondary SF (e.g., Palouš *et al.* 1990, Elmegreen 1994). Therefore, a comparison of the position of H I holes with H II regions should show some correlation, the H α emission being predominantly found along the rims of larger holes, as evidence for propagating SF. In the case of the smaller, young holes one would expect some to be still filled with diffuse H α emission coming from the ionized wind-blown cavities around the still present, young O and B stars.

Figure 10 shows an overlay of the H α map with the H I holes we detected in our survey (Sec. 4.1). The width of the rims of the H I holes is of course in reality larger than indicated in this plot. The subregions A–C will be discussed in more detail in the following section. As can be seen from this overlay, there is a prominent correlation of ongoing SF with the rims of the large H I shells which is in general agreement with the picture painted above. There is a strong tendency for H II regions to be situated on the rims of the H I shells (for some very nice examples see holes 8 and 35) and only in a very few cases diffuse H α emission does not seem to be related to an H I structure. Note that in one case only (hole 32), diffuse H α emission (partly) fills an H I hole.

Fig. 11 presents an overlay of the H α map with one contour of the H I surface brightness map (flux and primary beam corrected) plotted at 1.7×10^{21} atoms cm $^{-2}$. As can be seen from this overlay, H α emission is globally correlated with a high H I surface brightness (for an exception in the very north see below). Kennicutt (1989) used the disk instability relation of Toomre (1964) to show that, for spirals, there is a threshold density for star formation. For gas densities above the threshold, the star formation rate (SFR) is consistent with a Schmidt law. Near the threshold, the correlation breaks down, and one sees bursts of SF. Much below the threshold the SFR is extremely low. Skillman (1987) has observationally determined this threshold for dwarf galaxies to be on the order of 1×10^{21} atoms cm $^{-2}$ (at a resolution of 500 pc) which is in general agreement with our results (see also below). For a detailed discussion on the models that describe the relationship between gas, stars and star formation in irregular galaxies, the reader is referred to the paper by Hunter, Elmegreen & Baker (1998).

5.2. Analysis of individual H II regions

Using our high resolution H I data of IC 2574 we are now able to perform a much more detailed analysis and comparison of the properties of the H II regions with the H I features that are related to them. When making overlays of the H α map with the high resolution cube we found that in virtually all cases, single H II regions are correlated with distinct H I structures like clumps or clouds. We defined 40 regions where H α emission and H I clouds seem to be related. These regions are outlined by the squares in Figs. 12–14 which are meant to indicate which H II regions were lumped together. They are numbered 1' to

40' to avoid confusion with the numbers of the H I holes. Note, that sometimes two or more H α features belong to the same H I structure (like a clump) and were thus considered as one region (e.g., in region 40'), whereas in other cases individual H II regions were treated as separate entities (e.g., regions 11' and 12'). The H α fluxes of the regions tell us something about the current SF rate and the average number density of the ionized gas whereas the H I data yield approximations for the number density of neutral hydrogen, the velocity dispersion of the gas in that region and the thresholds for SF. Figs. 12–14 are enlargements of the regions A–C marked in Fig. 10 and display the regions defined by us. The position and size of the H I holes are plotted as well for comparison (please refer to Fig. 10 for cross referencing of the holes). The fluxes of the H II regions were taken from the observations of MH94. Table 5 summarizes the cross identifications between their and our regions. In this table we list the coordinates of the centers of our regions as well.

Table 6 summarizes the results of our analysis of the H II regions. The first column refers to the number of the H II region as defined in Figs. 12–14. From the fluxes published by MH94, we deduced the total flux $F(H\alpha)$ of our groups by simply summing their fluxes in the respective regions (column 2, Table 6). From this number, the H α luminosity can be calculated using $L(H\alpha) = 4\pi D^2 F(H\alpha)$, which, in the case of IC 2574 ($D=3.2$ Mpc), translates to $L(H\alpha) = 3.1 \times 10^{17} F(H\alpha)$ where $F(H\alpha)$ is given in $\text{ergs cm}^{-2} \text{s}^{-1}$ and $L(H\alpha)$ in units of solar luminosities $L_\odot$ ($1L_\odot = 3.85 \times 10^{33} \text{ ergs s}^{-1}$). The number of Lyman continuum photons per second is given by $N_{\text{Lyc}} = 2.82 \times 10^{45} L(H\alpha)$. Table 7 (see Devereux *et al.* 1997 and references

therein) summarizes the expected properties of high mass stars; column 1 gives the spectral type, column 2 the stellar mass, column 3 the bolometric stellar luminosity, column 4 the number of Lyman continuum photons per second, column 5 the H α luminosity $L(H\alpha)$, and column 6 the ratio of bolometric to H α luminosity. These equations and values can be used to estimate, for example, the equivalent number of O5 stars, for each region (see below).

Column 3 in Table 6 presents the volume density of the ionized free electrons over the area where we find H α emission (as defined by MH94). This was arrived at by first calculating the emission measure (EM) of each of their regions according to:

$$EM = 2.41 \times 10^3 T^{0.92} F(H\alpha) \text{ cm}^{-6} \text{ pc}$$

(Peimbert *et al.* 1975), where $F(H\alpha)$ is the H α flux in units of $\text{ergs cm}^{-2} \text{s}^{-1} \text{sterad}^{-1}$ and T is assumed to be the canonical value of 10^4 K. From that, the mean electron density for every single region published by MH94 was derived using

$$EM = n_e^2 d \text{ cm}^{-6} \text{ pc}$$

where n_e is the electron density and d the path length in pc through the H II region (assuming a spherical distribution of the H α emission). Since the derived values for n_e were very similar for a single region, we calculated the value for the electron density for each of our regions by taking an intensity-weighted average of the contributing MH94 regions. The average electron densities range from 0.5 to 2 cm^{-3} . This is, most likely, a lower limit for the actual electron density due to internal absorption within IC 2574. Note that in the above, effects due to the filling factor of the gas are ignored. The real densities in the H II regions are likely much higher.

In order to derive properties of the H I features that are physically related to the H II regions, we defined areas A around each of the H II regions with an effective radius given by $r_{\text{eff}} = \sqrt{A/\pi}$, where r_{eff} is given in pc (column 4). Although this is a somewhat subjective process, we found that slight variations in the size of the aperture didn't change the derived properties by more than 20 – 30%. Column 5 represents the total H I mass found within the aperture using the flux and primary beam corrected H I data. For every single region, we measured the velocity dispersion of the H I by taking spectra through the H I cube. Column 6 summarizes the measured velocity widths defined as the full width at half maximum (FWHM) from which the real velocity dispersion σ_v (assuming a Gaussian velocity distribution) can be calculated using $\Delta v = 2\sqrt{2\ln 2}\sigma_v = 2.355\sigma_v$. Interestingly, these values tend to be high where two or more shells overlap (e.g., the H II complexes 16, 31, 32, 33). These high dispersions are probably caused by blending along the line of sight of these shells.

Column 7 presents the average density of the neutral hydrogen which was calculated using

$$\rho_{\text{HI}} = M_{\text{HI}} / \left(\frac{4}{3} \pi r_{\text{eff}}^3 \right)$$

with M_{HI} and r_{eff} the values from columns 5 and 4, respectively. Values for ρ_{HI} range from about 1 cm^{-3} to values as high as 6 cm^{-3} . Note that these are only averages and that locally the H I density might be more enhanced. Based on the analysis by MCR94 and some crude mass modeling (Sec. 3.3) we derive an average density in the disk of $0.008 \text{ M}_{\odot} \text{ pc}^{-3} = 0.3 \text{ atom cm}^{-3}$. The regions in which star formation is currently active thus have an overdensity of an order of magnitude

(see Table 6).

The average surface density of the H I regions is shown in column 8. Note that virtually all values are larger than $1 \times 10^{21} \text{ cm}^{-2}$ which is in excellent agreement with the observational value given by Skillman (1987). In other words, SF is taking place if the projected H I surface density reaches values of about 10^{21} cm^{-2} not only for a resolution of 500 pc, as observed by Skillman but also for almost an order of magnitude improvement ($\approx 95 \text{ pc}$)! The highest surface density found is of order $2.5 \times 10^{21} \text{ cm}^{-2}$.

Finally Column 9 summarizes some remarks regarding the regions. Most of the H II regions are related to the rim of an H I shell. It should be noted at this point that in the course of checking for correlations between H α and H I emission, we detected 5 new tiny holes at the positions where there is H α emission. However, as these holes were not detected in our inspection of the H I cubes we didn't include them in the final catalogue which is thought to be complete for holes with radii larger than about 100 pc only (as discussed in Secs. 4.1 and 6.2).

We catalogued four of the H I features as “clumps” (regions 1', 2', 6' & 22'). By ‘clump’ we mean a well-defined H I region in 2 spatial dimensions as well as in velocity space. These clumps look like small, filled circles in position-velocity space. For those cases we asked ourselves whether these clumps are gravitationally confined or not. To answer that question, we calculated the virial masses of these objects using

$$M_{\text{vir}} = 250 \Delta v^2 r_{\text{eff}} \text{ M}_{\odot}$$

(Rohlf & Wilson 1996), where Δv is the full velocity width at half maximum (FWHM) in km s^{-1} and r_{eff} the radius of the clump in

pc. The virial masses of the four regions are: $M_{\text{vir}}(1') = 13 \times 10^6 M_{\odot}$, $M_{\text{vir}}(2') = 38 \times 10^6 M_{\odot}$, $M_{\text{vir}}(6') = 21 \times 10^6 M_{\odot}$ and $M_{\text{vir}}(22') = 7 \times 10^6 M_{\odot}$. As we see, the virial masses are all about one order of magnitude higher than the H I masses. So, the clumps are either not gravitationally bound or they are held together by as yet undetected material. One possibility is that they contain an order of magnitude more mass in the form of molecules. This will be hard to confirm, however, as dwarf galaxies have been notoriously difficult to detect in CO, the most abundant tracer for molecular gas (e.g., Sage *et al.* 1992). Dark Matter could play a role, but the mass density, including DM, near the above mentioned SF regions is of order $0.008 M_{\odot} \text{pc}^{-3}$ (see Sec. 3.3), insufficient by a wide margin.

If the observed velocity dispersion is not due to gravitation, the question remains, why *do* we see clumps? We can think of two more mechanisms, magnetic confinement or pressure. The former is unlikely to play a large role. Dwarf galaxies do not possess large scale magnetic fields (the exceptions being NGC 4449 and IC 10, e.g., Chyzy *et al.* 1998), any non-thermal radio continuum radiation being confined to the strongest regions of active star formation (Deeg *et al.* 1993).

We did not find any correlation of H α flux with the surrounding H I density. However, one wouldn't really expect any strong correlation of this kind. Not only is the H α emissivity of a star forming region strongly dependent on the age of the region (see, e.g., Gertsen & Icke 1997), but it is also dangerous to try to deduce actual physical properties of the SF regions from mean values, like average electron or H I densities, while ignoring spatial variations in their volume filling factors.

In some regions within IC 2574 (e.g., region 40'), we find high H I surface densities but only little H α emission indicating that possibly a major star formation burst is about to commence there and that therefore most of the gas is still neutral. With the help of the equations given above we can calculate how many stars are approximately needed to create the observed H α flux. In the case of region 40' we find a flux of $1.5 \times 10^{-14} \text{ergs cm}^{-2} \text{s}^{-1}$ and deduce a luminosity of $4.7 \times 10^3 L_{\odot}$ which is the equivalent of a few O7 stars only (see Table 7). Note, however, that there exist also regions where the situation is reversed and most of the H I is being ionized by the Lyman continuum photons of the most massive stars of the powerful H II regions. Region 34' is a prominent example (see also Fig. 11). From the H α flux of $F(H\alpha) \approx 1 \times 10^{-12} \text{ergs cm}^{-2} \text{s}^{-1}$ we derive an H α luminosity of $L(H\alpha) = 3.1 \times 10^5 L_{\odot}$ which corresponds to an equivalent of 22 O5 stars, in other words a major SF region.

6. Discussion

6.1. What created the H I holes?

The ISM of IC 2574 is dominated by expanding H I holes and shells. Similar structures have been found in other gas-rich dwarf galaxies (e.g., Holmberg II (PWBR92), IC 10 (Wilcots & Miller 1998) and in more massive spiral galaxies like our Galaxy (Heiles 1979), M 31 (Brinks & Bajaja 1988) and M 33 (Deul & den Hartog 1990). One of the major questions concerning these structures is what physical process or processes are responsible for their origin.

Heiles (1979), who was the first to recognize expanding H I shells in the Galaxy, concluded that 'many of the H I structures

[...] were probably produced by large amounts of energy injected directly to the interstellar medium'. One obvious mechanism is the after effects of massive star formation: strong stellar winds and supernova explosions (see the review by Tenorio-Tagle and Bodenheimer 1988). Detailed numerical modeling by various authors (e.g., Palouš, Franco & Tenorio-Tagle 1990, Silich *et al.* 1996, Palouš, Tenorio-Tagle & Franco 1994) has strengthened confidence in this explanation, as has the work by Oey & Clarke (1997) who in a (semi-)analytical way managed to predict the H I shell size distribution in the SMC based on the observed H α luminosity function.

Having said that, there are potentially problems with this interpretation, as mentioned by Radice, Salzer & Westpfahl (1995) and Rhode, Salzer & Westpfahl (1999). Looking for remnant stellar clusters at the centers of several large holes in Holmberg II they failed to find a population of A and F main sequence stars. Moreover, it is difficult to imagine that the largest H I shells which are located beyond the optical galaxy should have been created by star formation (an argument which holds equally well for some of the holes in IC 2574). This is corroborated by ultraviolet imaging of Holmberg II by Stewart (1999) who finds no FUV emission from the outer regions of Holmberg II.

It should be noted, though, that the non-detection of population A and F stars does not necessarily mean that none of the H I are created by SF. In dwarf galaxies, in the absence of shear, H I shells can reach large dimensions and grow so old that no obvious bright cluster remnant will remain visible in the center (Efremov, Elmegreen & Hodge). Some of the observed supershells are so old that the brightest members of stellar cluster

which created the hole may have evolved off the main sequence and dispersed (assuming a similar IMF as in the Galaxy). This may make a detection of old clusters difficult. Such a case (LMC 4/Constellation III) is described by Efremov & Elmegreen (1998).

Another potential problem with the conventional picture for the origin of the holes is that for the largest ones the energy requirements seem to surpass the amount that can possibly be released into the ambient ISM via stellar winds and supernova explosions. The giant shell in M 101 (van der Hulst & Sancisi 1988) is just one example of such a structure. Tenorio-Tagle (1987) proposed as an alternative the infall of high velocity clouds which could deposit an amount of energy of 10^{52} to 10^{54} ergs per collision (see the review by Tenorio-Tagle & Bodenheimer 1988 for further references). Even though this mechanism seems a viable explanation for at least some holes, it is unlikely that it plays a dominant role. Certainly it can't be as important in dwarf galaxies because of their much smaller collisional cross section.

Recently, an alternative explanation for the creation of supergiant H I shells in galaxies has been proposed, namely GRBs or Gamma Ray Bursters (Efremov, Elmegreen & Hodge 1998, Loeb & Perna 1998). Even though the basic physics of GRB are not yet known the picture to emerge is that they are highly energetic events (of order 10^{54} ergs) that release huge amounts of energy in a very short period of time (order of a few seconds).

Although we recognize that perhaps not all holes can be explained by stellar winds and supernova explosions, we feel that there is sufficient evidence for the conventional interpretation to warrant an analysis in those terms, which is what we pretend in the rest of this

paper.

6.2. Statistical properties of the holes: a comparison with other galaxies

For the first time we are now in a position to address an important question regarding the creation and evolution of hole-like structures in the interstellar medium: how are their properties related to their environment, i.e., the type of host galaxy. In what follows we will compare the observed and derived H I hole properties of IC 2574 with those found in M 31 (Brinks & Bajaja 1986), an example of a massive spiral galaxy similar to our own, M 33 (Deul & den Hartog 1990), a less massive spiral and Ho II (PWBR92), another dwarf galaxy in the same group of galaxies, yet four times less massive than IC 2574. In other words, the sequence M 31 – M 33 – IC 2574 – Ho II spans a large range of different Hubble types from massive spirals to low-mass dwarfs.

Despite the fact that a similar study, and at considerably higher sensitivity, is now available for the SMC (Staveley-Smith *et al.* 1997), we decided not to include their results. The main reason being that the linear scales (or spatial frequencies) sampled by the ATNF hardly overlap with those observed in the galaxies listed above. The linear resolution, at 28 pc, is almost four times higher than, for example, the VLA maps on IC 2574. At the other end of the spectrum, because of the lack of short spacing information, structures larger than a few hundred parsec will have been missed. Moreover, the SMC is a very disturbed system, being torn apart by tidal forces due to interactions with the LMC and the Galaxy.

Limiting the comparison to the four galaxies listed above has some further advantages.

The linear resolutions are very similar, as are the velocity resolutions with which they have been observed (see Tab. ?? for a summary). In addition, all four galaxies were examined in more or less the same fashion, one of the authors (EB) having taken part in the analysis of three of the four objects. We are aware that the results for the four galaxies suffer partially from low statistics and incompleteness due to personal bias and observational constraints (such as the beamsizes). However, we feel that these effects, to first order, affect a comparison in a similar way and that it is valid to try to find global trends as a function of Hubble type.

For the sake of brevity, rather than first show the statistical properties of the holes in IC 2574 separately, we immediately go to a comparison with the other three objects. In Figs. 15–19 we compare the distributions of swept-up matter, the energies, diameters, expansion velocities and ages of the holes in the 4 galaxies. In each graph we plot in the form of histograms the relative number of holes, in percent, in order to make a direct comparison possible. In each histogram, the data for the individual objects are binned in the same way. To improve the presentation, however, the respective bins were slightly shifted when plotting the results.

Fig. 15 shows an overlay of the indicative masses of the H I holes, binned logarithmically. The values range from typically 10^4 to $5 \times 10^6 M_\odot$. Even though a two-sample Kolmogorov–Smirnov test rejects the hypothesis of pairwise equality of the distributions, a casual inspection shows that the indicative masses span roughly the same range. The mean masses (in $10^5 M_\odot$) for the dwarf galaxies (Ho II: 10.5, IC 2574: 8.4) are slightly higher as compared to the spirals (M 33: 4.1,

M 31: 3.2). This is somewhat surprising if one realizes that the volume density in the plane of a spiral galaxy is typically higher by a factor of three as compared to that found in a dwarf galaxy. H I column densities have a much smaller variation from object to object, though (see Tab. ??) which probably explains why the swept up matter works out to be more or less the same. As a warning, it should be born in mind that, in order to determine the H I densities we smeared out any fine scale structure, potentially adding somewhat to the errors.

Fig. 16 is similar to Fig. 15, showing instead the distribution of the logarithm of the energies needed to create the observed H I holes (under the assumption of them being created by supernova explosions). The derived energies range from 10^{50} to 10^{53} ergs. We find the important result that the distributions are identical, at least to first order (mean energies (10^{52} ergs): Ho II: 1.4, IC 2574: 1.1, M 33: 1.6, M 31: 1.5). A two-sample Kolmogorov–Smirnov test reveals that there is no significant pairwise difference at the 95% level between the four energy distributions. From this we conclude that regardless of Hubble type of the parent galaxy, the star clusters (or whatever other process created the H I holes) deposit more or less the same amount of energy into the ISM.

Fig. 17 shows an overlay of the size distribution of the holes found in the four galaxies. In this and the following plots the bins are on a linear scale. Note that there is a clear difference with Hubble type! The size distribution for holes in M 31 and M 33 cuts off sharply near 600 pc. In contrast, holes in IC 2574 and Ho II reach sizes of 1200 to 1500 pc, respectively. The lack of holes with sizes smaller than ~ 100 pc is due to our resolution limit.

The fact that the holes get larger for “later” Hubble types (i.e., galaxies with lower mass) can be understood as follows. Smaller galaxies have lower masses and their mass surface density in the disk is lower as well. The one-dimensional velocity dispersion varies little, if any, among galaxies. We find in the four objects under scrutiny, taking into account an uncertainty of $\pm 1 \text{ km s}^{-1}$, a range of 6–9 km s^{-1} . This, combined with the lower mass surface density, leads to dwarf galaxies having much thicker disks with aspect ratios of 1:10 rather than 1:100. Hence, for the same amount of energy deposited, as argued above, an H I shell can grow much larger, because of a lower gravitational potential and a lower ambient density. Moreover, in dwarf galaxies shells are prevented from breaking out of the disk because it is thicker (see Sec. 3.3).

The age distribution of the holes (Fig. 18) also shows a sequence with Hubble type. The mean values for the ages of the holes are: 10×10^6 years (M 31); 12×10^6 years (M 33); 23×10^6 years (IC 2574) and 62×10^6 years (Ho II). In the spirals no holes are found which are older than 30 Myr whereas the distribution in dwarfs is flatter and spans an age range of up to 120 Myr. This is not unexpected since in more massive galaxies shear and mixing due to the passage of density waves tend to destroy holes. In dwarfs, unless a shell is hit by a neighboring explosion, and subsequently expanding structure, a feature can potentially persist for a long time, until the rim disperses and mixes with the surrounding ISM.

Finally, the distribution of the expansion velocities is presented in Fig. 19. The predominantly low values for the dwarfs (especially Ho II) reflect, again, the fact that we see many large holes in the dwarfs which have

almost stalled. A word of caution should be added as the velocity resolution of the IC 2574 and Ho II observations was a factor of about 3 better than those of M 31. Very young (i.e., smaller than ~ 100 pc) and fast expanding holes are missed, of course, but this holds true for all galaxies in the sample.

Figure 20 displays an alternative way of looking at the properties of the H I holes on a galaxy-wide scale. It shows the expansion velocities of the shells in the four nearby galaxies plotted as a function of their diameter (crosses). Based on the hydrodynamical models for expanding shells (Chevalier 1974), we plotted lines of constant energy for the expanding structures in Fig. 20 (assuming a constant density of the ambient medium of 0.15 cm^{-2}). The plotted curves represent energies at a level of 10^{49} , 10^{50} , 10^{51} and 10^{52} ergs, respectively (from left to right). The evolution of an individual hole of a given energy is along an equi-energy line from the top of the plot downwards. In addition, lines of equal ages are shown in the figure, ranging from 10^7 years (steepest slope) via 5×10^7 years to 10^8 years.

There is a marked difference between the two spiral galaxies and the dwarfs. The H I shells in M 31 and M 33 are smaller and the range of expansion velocities is larger. In fact, when looking at the objects in order of decreasing mass (M 31 – M 33 – IC 2574 – Ho II) there seems to be a gradually shifting pattern. Whereas in the larger spirals there is an absence of older holes (older than a few $\times 10^7$ years, holes in IC 2574 range between 5×10^7 and 10^8 years and holes in Ho II cluster near 10^8 years

This suggests an alternative explanation for why we don't see younger holes in IC 2574 and Ho II. The SF histories of these dwarf

galaxies could be quite different from the spirals. In the latter, the global SF rate is, to first order, constant with time. Therefore, there is always a population of young holes. In contrast, in gas-rich dwarf irregulars, SF occurs in bursts which transforms them into an H II or blue compact dwarf (BCD) galaxy. It might be that IC 2574 and Ho II have undergone a recent burst, perhaps triggered by a close passage with another member of the M 81 group, and are now in their post-starburst phase and that this is why we don't see many young holes. Our data alone don't allow us to choose between the two possibilities.

Oey & Clarke (1997), using the standard, adiabatic shell evolution predict the size distribution for the populations of OB superbubbles. In their paper they compare their predictions with observations of the SMC and the three same galaxies as we compared our IC 2574 data with, namely M31, M33, and Ho II. Without going into detail, their prediction fits the observations for the SMC surprisingly well. This despite the fact that they ignore the selection effect (missing short spacings) which renders the SMC maps insensitive to large structures which should dominate the appearance of dwarf galaxies. Also, they emphasize stalled shells, i.e., objects which have reached their largest diameter. This is in contradiction with the fact that the SMC shells, especially the smaller ones, are all found to be rapidly expanding. Their approach works less well for the other objects. They ascribe this to the data being incomplete, not only when considering the smallest detectable shells, but overall, due to the reduced sensitivity as compared to the SMC data. In that respect, our IC 2574 are not better than the maps on M31, M33 or Ho II. Hence, a thorough test of Oey &

Clarke’s predictions will have to await higher resolution and higher sensitivity data on the nearest (dwarf) galaxies.

6.3. The intrinsic shape of the shells

One of the results from our analysis is that over 90% of the holes are well approximated by a circular shape (the few exceptions are discussed in Sec. 4.2). This is a major difference with other galaxies studied thus far in similar detail (i.e., Ho II, M 31, M 33, M 101, NGC 6946) which tend to have much more elliptical holes. One attempt to understand this is in terms of projection effects (different inclinations for the galaxies observed thus far). If the shells are intrinsically triaxial, their projected shape on the sky will invariably be an ellipse with an ellipticity which varies as a function of the intrinsic axial ratios, the inclination of the galaxy, *and* the position angle of the major axis of the hole within the plane of the galaxy. This should lead to a wide range of ellipticities. The fact that we don’t see this in IC 2574 can only mean that the shells in this galaxy are *intrinsically* spherical.

This can be understood by considering the shapes of the rotation curves of the galaxies. IC 2574 is the only object which exhibits a rising rotation curve throughout, as is typical for dwarf galaxies. As the rotation curve does not turn over, leading to a flat part dominated by differential rotation, there is no shear to distort and ultimately destroy any large scale structures such as H I shells. For a more complete discussion on this topic, see Silich *et al.* (1996) and Palouš *et al.* (1990). Indeed, PWBR92 estimate that the amount of shear over 1 or 2 kpc in radius, once the flat part of the rotation curve is reached in Ho II, is sufficient to stretch the holes to an axial ratio of 0.5 over a timescale of 10^8 years.

IC 2574, however, is the only object studied to date which shows solid body rotation almost throughout the disk (MCR94) which ensures that holes stay circular until long after their creation.

There is one fly in the ointment. Why should the shells in IC 2574 be intrinsically spherical? The expansion rate perpendicular to the disk should be different from that in the disk. In the direction perpendicular to the disk, the volume density rapidly decreases, favoring faster expansion in that direction. But on the other hand, the gas which moves upwards starts to feel the gravitational pull from the mass surface density below it, slowing it down. It thus seems as if nature conspires to create, at least in this particular object, holes which are near spherical in shape.

Based on the absence of elliptical holes, an interesting conclusion can be drawn concerning magnetic fields. Usually, the influence of magnetic fields is neglected in numerical studies of expanding bubbles in the ISM. However, a few analytical approaches have been put forward in recent years to investigate the effects of the evolution of shell-like structures in the presence of magnetic fields (e.g., Tomisaka 1991). The results of these investigations are that if global magnetic fields are present, this would inhibit the expansion of a supernova driven bubble in directions perpendicular to the field lines. The expansion along the field lines would not be affected, thus leading to elliptical, elongated structures. The fact that we observe only spherical holes in IC 2574 shows that either magnetic fields play a minor role or that the magnetic fields in IC 2574 are really weak.

7. Summary

High resolution H I and H α imaging using the VLA and the Calar Alto observatory of the dwarf galaxy IC 2574 have been presented in this paper. The main results and conclusions are:

1. IC 2574 is a gas-rich dwarf galaxy which shows a stunning amount of detail in the form of H I shells and holes in its interstellar medium. These features are similar to those found in the Galaxy, the Magellanic Clouds, M 31, M 33, M 101 and NGC 6946, IC 10 and Ho II. From the global H I profile we derive a total H I mass of $9.0 \times 10^8 M_\odot$.

2. Using two independent approaches, the thickness of the H I layer of IC 2574 has been derived to be ≈ 350 pc. This implies that IC 2574, as is the case for Ho II, is a 'thick' dwarf galaxy (compared to more massive spirals) which can be understood in terms of its lower gravitational potential.

3. Large scale structures emptied of H I (believed to be expanding H I shells) leave roughly circular footprints, H I holes. They are prominent in the high resolution VLA channel maps of IC 2574. In total, 48 H I holes have been catalogued which are distributed across the entire galaxy. The diameters of the holes range between 100 pc and 1200 pc. The lower limit of the sizes is due to the limit of our spatial resolution. Most of the holes are found to be expanding with typical expansion velocities ranging between 8 and 12 km s $^{-1}$. The indicative ages of the H I holes range between 10 and 60×10^6 years. Typical energies that are required to create the observed features are of the order of $10\text{--}100 \times 10^{50}$ ergs assuming that they were created by the combined effects of stellar winds and supernova explosions. The

H I masses that were present at the position of the holes before creation (and that might now be present on the rim of the shell) are of the order of $10\text{--}100 \times 10^4 M_\odot$.

4. In addition to the well defined holes, some large scale (> 1000 pc), coherent features are visible in the channel maps. They might be the remainder of an older shell population.

5. The radial expansion of the holes, the indicative ages and the energy requirements for their formation suggest that they have been created by the combined effects of stellar winds and multiple supernova explosions. Despite their large dimensions, the energy requirements are not extreme and infall of material such as clouds is not needed to explain the observed features. However, alternative mechanisms cannot be ruled out.

6. The shells can grow to these large dimensions because of several conditions which are fulfilled in dwarf galaxies. The volume density in the plane is low, which facilitates expansion. In the direction perpendicular to the disk, the gravitational pull is smaller than in a massive spiral. Also, because of the thick H I layer, shells are easily contained and unlikely to blow-out. Lastly, solid body rotation and a lack of spiral density waves prevent holes from being rapidly destroyed.

7. A comparison with other galaxies studied in similar detail so far (M 31, M 33 and Ho II) shows that the size distribution of H I holes found in a galaxy is related to its Hubble type in the following way. The size of the largest H I shells is inversely proportional to the global gravitational potential (and hence mass surface density). The shells in dwarf galaxies show lower expansion velocities, most likely due to the fact that the shells are preserved much longer due to the absence

of shear (differential rotation). The energies needed to create these structures, though, are found to be roughly the same for all types of galaxies. If we follow the conventional view that star forming regions are responsible for the structures found in H I, the energy output of a typical star forming region does not seem to be related to the host galaxy.

8. The fact that virtually all holes in IC 2574 are circular in shape is a major difference with the other galaxies studied in comparable detail and can be explained, again, by the absence of shear and spiral density waves in IC 2574. This finding also indicates that magnetic fields play only a minor role in shaping expanding holes or that the magnetic fields in IC 2574 are weak.

9. Current star formation, as traced by H α emission, is predominantly found along the rims of the larger H I-holes, indicating propagating star formation. A detailed comparison of the H α and the H I properties yields that, on a global scale, star formation is only present if the H I surface density reaches values larger than 10^{21} cm^{-2} (on a linear scale of $\approx 95 \text{ pc}$).

In summary, our results on IC 2574 confirm the findings by PWBR92 and show that Ho II is not a special case. Our analysis, and the comparison with other galaxies studied in similar detail suggest that there is a trend of hole characteristics with Hubble type (or mass). If the conclusion that the energy output of a SF region is independent of the host galaxy holds then it follows that there is a lower mass limit for the formation of dwarf galaxies. In other words, if the mass of a galaxy falls below a certain threshold, the first burst of star formation will tear the whole galaxy apart (see also, e.g., Puche & Westpfahl 1994). Mac Low & Ferrara (1998) pre-

dict that this lower mass limit will be near $10^6 M_{\odot}$, which is well below the mass of the objects dealt with in this paper.

The authors are indebted to Mordecai-Marc Mac Low for providing them with a model data cube of a turbulent ISM in advance of publication and to Johan van Horebeek for valuable advice regarding some of the statistical techniques employed. FW acknowledges the 'Deutsche Forschungsgemeinschaft (DFG)' for the award of a stipendium in the framework of the Graduiertenkolleg on "The Magellanic Clouds and other Dwarf Galaxies" and would like to thank Uli Klein for continuous support and Neb Duric for fruitful discussions. FW appreciates the help of the staff of the Array Operation Center of the NRAO during his stay in Socorro, New Mexico and also wants to thank the staff of the Calar Alto observatory in Spain for excellent working conditions. FW is grateful to Jan Palouš for illuminating discussions regarding the numerical simulations of holes and shells during his stays at the Academy of Science in Prague, Czech Republic. EB acknowledges a grant awarded by CONACyT (grant number 0460P-E) and is grateful for the support by the Graduiertenkolleg without which this collaborative effort would not have been possible. Last but not least we would like to thank Klaas de Boer for carefully reading the manuscript and an anonymous referee for useful suggestions. This research has made use of the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, Caltech, under contract with the National Aeronautics and Space Administration (NASA), NASA's Astrophysical Data System Abstract Service (ADS), and NASA's SkyView.

REFERENCES

- Arsenault, R., & Roy, J.R. 1986, *AJ*, 92, 567
- Baars, J.W.M., Genzel, R., Pauliny-Toth, I.I.K., & Witzel, A. 1977, *A&A*, 61, 99
- Bottema, R., Shostak, G.S., & van der Kruit, P.C. 1986, *A&A*, 167, 34
- Briggs, D. 1995, Ph.D. thesis, New Mexico Institute of Mining and Technology
- Brinks, E. 1981, *A&A*, 95, L1
- Brinks, E., & Bajaja, E. 1986, *A&A*, 169, 14
- Chevalier, R. A. 1974, *ApJ*, 188, 501
- Chyzy, K.T., Kohle, S., Beck, R., Klein, U. & Urbanik, M. 1998, in *Proceedings of the Bonn/Bochum-Graduiertenkolleg Workshop: The Magellanic Clouds and Other Dwarf Galaxies*, ed. T. Richtler, & J. M. Braun (Shaker Verlag:Bonn), p. 237
- Clark, B.G. 1980, *A&A*, 89, 377
- Cox, D.P., & Smith, B.W. 1974, *ApJ*, 189, L105
- Deeg, H.-J., Brinks, E., Duric, N., Klein, U., & Skillman, E. 1993, *ApJ*, 410, 626
- Deul, E.R., & den Hartog, R.H. 1990, *A&A*, 229, 362
- Devereux, N.A., Duric, N., & Scowen, P.A., 1997, *AJ*, 113, 236
- de Vaucouleurs, G., de Vaucouleurs, A., Corwin, H.G., Buta, R.J., Paturel, G., & Fouqué, P. 1991, *Third Reference Catalogue of Bright Galaxies* (Springer:New York)(RC3)
- Dressel, L.L., Condon, J.J. 1976, *A&AS*, 31, 187
- Drissen, L., Roy, J.-R., & Moffat, A.F.J. 1993, *AJ*, 106, 1460
- Efremov, Y.N., & Elmegreen, B.G. 1998, *MNRAS*, 299, 643
- Efremov, Y.N., Elmegreen, B.G., & Hodge, P.W. 1998, *ApJ*, 501, L163
- Elmegreen, B.G. 1994, *ApJ*, 427, 384
- Field, G.B., Goldsmith, D.W., & Habing, H.J. 1969, *ApJ*, 155, L149
- Gerritsen, J.P.E., Icke, V. 1997, *A&A*, 325, 972
- Heiles, C. 1979, *ApJ*, 229, 533
- Heiles, C. 1984, *ApJS*, 55, 585
- Heiles, C. 1990, *ApJ*, 354, 483
- Högbom, J.A. 1974, *A&AS*, 248, 1
- Huchtmeier, W.K., & Richter, O.G. 1988, *A&A*, 203, 237
- Hunter, D.A., Elmegreen, B.G., Baker, A.L. 1998, *ApJ*, 493, 595
- Jörsäter, S., & van Moorsel, G.A. 1995, *AJ*, 110, 2037
- Kamphuis, J.J. 1993, Ph.D. thesis, University of Groningen
- Kellmann, S.A. 1972, *ApJ*, 175, 353
- Kennicutt, R.C. 1988, *ApJ*, 334, 144
- Kennicutt, R.C. 1989, *ApJ*, 344, 685
- Kim, S., Staveley-Smith, L., Dopita, M.A., Freeman, K.C., Sault, R.J., Kesteven, M.J., & McConnell, D. 1999, *ApJ*, 503, 729
- Loeb, A., & Perna, R. 1998, *ApJ*, 503, L35
- Martimbeau, N., Carignan, C., & Roy, J.-R. 1994, *AJ*, 427, 656 (MCR94)
- McCray, R., & Snow, T.P. Jr. 1979, *ARA&A*, 17, 213
- Mac Low, M.-M., & Ferrara, A. 1998, *ApJ*, 513, in press

- Mac Low, M.-M., Klessen, R.S., Burkert, A., & Smith, M.D. 1998, *Phys. Rev. Lett.*, 80, 2754
- McKee, C.F. & Ostriker, J.P. 1977, *ApJ*, 218, 148
- Miller, B.W., & Hodge, P. 1994, *ApJ*, 427, 656 (MH94)
- Miller, B.W., & Hodge, P. 1996, *ApJ*, 458, 467
- Norman, C.A., & Ikeuchi, S. 1989. *ApJ*, 345, 372
- Oey, M.S., & Clarke, C.J. 1997, *MNRAS*, 289, 570
- Palouš, J., Franco, J., & Tenorio-Tagle, G. 1990, *A&A*, 227, 175
- Palouš, J., Tenorio-Tagle, G., & Franco, J. 1994, *MNRAS*, 270, 75
- Peimbert, M., Rayo, J.F., & Torres-Peimbert, S. 1975, *RMx A*, 1, 289
- Puche, D., Westpfahl, D., Brinks, E., & Roy, J.-R. 1992, *AJ*, 103, 1841 (PWBR92)
- Puche, D., & Westpfahl, D. 1994, in *ESO/OHP Conference and Workshop Proceedings No. 49*, ed. G. Meylan, & P. Prugniel (ESO:Garching), 273
- Radice, L.A., Salzer, J.J., & Westpfahl, D.J. 1995, *BAAS*, 186, 4908
- Rhode, K.L., Salzer, J.J., & Westpfahl, D.J. 1999, *AJ*, submitted.
- Rohlfs, K., & Wilson, T.L. 1996, *Tools of Radio Astronomy* (Springer Verlag:Berlin)
- Rots, A.H. 1980, *A&AS*, 41, 189
- Sage, L.J., Salzer, J.J., Loose, H.-H., & Henkel, C. 1992, *A&A*, 265, 19
- Seielstad, G.A., & Wright, M.C.H. 1973, *ApJ*, 184, 343
- Shostak, G.S., & Skillman, E.D. 1989, *A&A*, 214, 33
- Silich, S.A., Franco, J., Palouš, J., & Tenorio-Tagle, G. 1996, *ApJ*, 468, 722
- Skillman, E.D. 1987, in *NASA Conf. Pub. CP-2466: Star Formation in Galaxies*, ed. C.J. Lonsdale Persson (NASA:Washington), 263
- Staveley-Smith, L., Davies, R.D., & Kinman, T.D. 1992, *MNRAS*, 258, 334
- Staveley-Smith, L., Sault, R.J., Hatzidimitrou, D., Kesteven, M.J., & McConnell, D. 1997, *MNRAS*, 289, 225
- Stewart, S. G., et al. 1999, *ApJ*, submitted.
- Tenorio-Tagle, G., Franco, J., Bodenheimer, P., & Różyczka, M. 1987, *A&A*, 179, 219
- Tenorio-Tagle, G., & Bodenheimer, P. 1988, *ARA&A*, 26, 145
- Tikhonov, N.A., Karachentsev, I.D., Bilkina, B.I., & Georgiev, T.B. 1991, *A&AS*, 89, 1
- Tomisaka 1991, in *IAU Symposium No. 144: The Interstellar Disk-Halo Connection in Galaxies*, ed. H. Bloemen (Kluwer:Dordrecht), 407
- Tomita, A., Ohta, K., Nakanishi, K., Takeuchi, T.T., & Saito, M. 1998, *AJ*, submitted
- Toomre, A. 1964, *ApJ*, 139, 1217
- van der Hulst, J. M., & Sancisi, R. 1988, *AJ*, 95, 1354
- van der Hulst, J. M. 1996, in *ASP Conf. Ser. Vol. 106: The Minnesota Lectures on Extragalactic Neutral Hydrogen*, ed. E. D. Skillman (ASP:San Francisco), 47
- van der Kruit, P.C. 1981, *A&A*, 99, 298
- Walter, F., Brinks, E., Duric, N., & Klein, U. 1997, *AJ*, 113, 2031

Walter, F., Brinks, E., Duric, N., Kerp, J., & Klein, U. 1998, *ApJ*, 502, L143

Wilcots, E.M., & Miller, B.W. 1998, *AJ*, 116, 2363

Fig. 1.— R-band CCD-image of IC 2574 obtained with the Calar Alto 2.2-m telescope.

Fig. 2.— Continuum-subtracted $H\alpha$ image of IC 2574 obtained with the Calar Alto 2.2-m telescope. The total exposure time is 45 minutes. The brightest stars that were not fully subtracted by removing the continuum were blanked.

Fig. 3.— Mosaic of every second channel map of IC 2574 in which line emission was detected. The velocity of each channel in km s^{-1} is plotted in the right hand corner. Greyscale range from 0 (white) to 6 mJy beam^{-1} (black). Note that the beamsize ($6.4'' \times 5.9''$ as indicated in the box of the top left hand panel) is smaller than the resolution of this plot. Coordinates are with respect to B1950.0 equinox.

Fig. 4.— Global H I profile for IC 2574 (squares) obtained by integrating the channel maps over the area of the source after subtraction of the continuum background. The data have been flux corrected (see text) and are corrected for primary beam attenuation. The Flux F is given in Jy. The channel spacing is 2.58 km s^{-1} . For comparison, the profiles published by Rots (1980) (full line) and MCR94 (dotted line) are also plotted. The arrow indicates the velocity range affected by contamination due to galactic emission.

Fig. 5.— H I surface brightness map of IC 2574. The greyscale is a linear representation of the H I surface brightness ($0 - 2406 \text{ K km s}^{-1}$). The beamsize ($6.4'' \times 5.9''$) is indicated in the lower left.

Fig. 6.— Isovelocity contours of IC 2574 of the $30''$ convolved cube. The velocities are

This 2-column preprint was prepared with the AAS L^AT_EX macros v4.0.

given in km s^{-1} . The greyscale image is a linear representation of the H I surface brightness map at full resolution. A circular $30''$ beam is plotted in the lower left hand corner.

Fig. 7.— Velocity dispersion (second moment) map of IC 2574. The greyscale is a linear representation of the velocity dispersion ranging between 4 km s^{-1} (white) and 13 km s^{-1} (black). The beamsize ($6.4'' \times 5.9''$) is indicated in the lower left.

Fig. 8.— Typical examples of position velocity diagrams. a) pV diagram of a Type 3 hole (hole number 21). b) rV diagram of the same hole. c) pV diagram of a Type 1 hole (number 30) and d) pV diagram of a Type 2 hole (number 33).

Fig. 9.— Location of the H I holes in IC 2574 (with numbers). The greyscale map is a linear representation of the H I surface brightness map. For the largest holes (13, 23, and 26), see text.

Fig. 10.— $\text{H}\alpha$ image of IC 2574 overlaid with the discovered holes in its neutral interstellar medium (with numbers). Regions A–C are enlarged in Figs. 12–14, respectively. Note that almost all $\text{H}\alpha$ emission in IC 2574 is located on the rims of H I shells (see text).

Fig. 11.— $\text{H}\alpha$ image of IC 2574 overlaid with one contour of the H I surface brightness map at $1.7 \times 10^{21} \text{ atoms cm}^{-2}$. Note that regions where we find $\text{H}\alpha$ emission coincide with high H I surface brightness.

Fig. 12.— Enlargement of region A from Fig. 10. The circles indicate where we detected the H I holes (for identification of the H I holes, please refer to Fig. 10). The boxes

show how we defined groups of H II regions within IC 2574 (with sequence numbers).

Fig. 13.— Enlargement of region B from Fig. 10. The circles indicate where we detected the H I holes (for identification of the H I holes, please refer to Fig. 10). The boxes show how we defined groups of H II regions within IC 2574 (with sequence numbers).

Fig. 14.— Enlargement of region C from Fig. 10. The circles indicate where we detected the H I holes (for identification of the H I holes, please refer to Fig. 10). The boxes show how we defined groups of H II regions within IC 2574 (with sequence numbers).

Fig. 15.— Comparison of the relative distribution, in percentage, of the indicative masses of the H I holes in Ho II, IC 2574, M 33 and M 31.

Fig. 16.— Comparison of the relative distribution, in percentage, of the energies required to produce the H I holes in Ho II, IC 2574, M 33 and M 31.

Fig. 17.— Comparison of the relative distribution, in percentage, of the diameters of the H I holes in Ho II, IC 2574, M 33 and M 31.

Fig. 18.— Comparison of the relative distribution, in percentage, of the ages of the H I holes in Ho II, IC 2574, M 33 and M 31.

Fig. 19.— Comparison of the relative distribution, in percentage, of the expansion velocities of the H I holes in Ho II, IC 2574, M 33 and M 31.

Fig. 20.— The relation between expansion velocity and diameter in each of the four galaxies. Crosses represent observed values. The plotted curves are lines of constant energy

(based on Chevalier 1974) at levels of 10^{49} , 10^{50} , 10^{51} and 10^{52} ergs (from left to right). The straight lines correspond to constant age and are drawn at 10^7 years (steepest slope), 5×10^7 years and 10^8 years.

Table 1: General information on IC 2574

Object	IC 2574
Other names	UGC 5666, DDO 81, VII Zw 330
Right ascension (1950.0) ^a	10 ^h 24 ^m 41 ^s .3
Declination (1950.0) ^a	68° 40′ 18″
Adopted distance ^b	3.2 Mpc
Inclination ^c	77±3°
Scale (1″ corresponds to)	15.5 pc
Apparent magnitude (B) ^d	10.8 mag
Corrected distance modulus	m _B -M _B =27.6
Corrected absolute magnitude (B)	-16.8 mag
Blue luminosity L _B	7.8×10 ⁸ L _{B⊙}
HI mass M _{HI} ^e	9.0×10 ⁸ M _⊙
HI mass to blue light ratio M _{HI} /L _B	1.15 M _⊙ /L _{B⊙}

^aDressel & Condon 1976

^bsee discussion in Puche *et al.* 1992

^cMartimbeau *et al.* 1994

^dde Vaucouleurs *et al.* 1991 (RC3)

^ethis study

Table 2: Setup of the VLA during the observations

VLA configuration	B	C	D
baselines	1 – 54 k λ (0.21 – 11.4 km)	0.34 – 16 k λ (0.073 – 3.4 km)	0.166 – 4.9 k λ (0.035 – 1.03 km)
date of observation	1992 Jan. 18 1992 Jan. 19	1992 Apr. 20	1991 Mar. 03 1992 Sep. 03
total time on source	647 min	133 min	245 min
total bandwidth		1.56 MHz	
No. of channels		128	
velocity resolution		2.58 km s ⁻¹	
central velocity		38 km s ⁻¹	
angular resolution ^a		6.4 $\times$ 5.9 arcsec	
linear resolution ^a		100 $\times$ 91 pc	
rms noise ^a		0.7 mJy $\times$ beam ⁻¹ 11 Kelvin	

^abased on ROBUST data cube; see text

Table 3: Observed properties of the H I holes

#	α (1950.0)	δ (1950.0)	v_{hel} (km s^{-1})	d (pc)	v_{exp} (km s^{-1})	n_{HI} (cm^{-3})	type
1	10 23 34.5	68 36 26	-11	310	8	0.13	3
2	10 23 38.0	68 36 43	-1	310	5	0.13	3
3	10 23 48.0	68 36 58	5	976	17	0.15	3
4	10 23 52.5	68 40 14	28	1240×1116,42°	14	0.05	2
5	10 23 59.0	68 34 07	-1	775	9	0.1	3
6	10 24 00.5	68 42 08	53	325	8	0.04	3
7	10 24 02.5	68 36 50	2	1162	—	0.16	1
8	10 24 04.6	68 39 57	33	372	15	0.08	3
9	10 24 05.5	68 37 59	17	418	7	0.16	3
10	10 24 06.5	68 36 40	7	418	—	0.15	1
11	10 24 12.0	68 42 33	43	697	8	0.04	3
12	10 24 15.0	68 37 12	15	325	—	0.16	1
13	10 24 15.0	68 39 13	25	1922	—	0.16	1
14	10 24 17.0	68 38 11	20	325	—	0.17	1
15	10 24 23.5	68 37 20	25	697	6	0.16	3
16	10 24 26.5	68 37 57	25	325	8	0.16	3
17	10 24 26.5	68 43 33	53	465	13	0.08	3
18	10 24 28.0	68 38 43	28	434	11	0.17	3
19	10 24 31.5	68 42 58	36	232	8	0.07	3
20	10 24 42.0	68 40 18	33	837	—	0.16	1
21	10 24 35.0	68 39 48	38	604	18	0.17	3
22	10 24 40.5	68 39 35	38	976×558,78°	13	0.17	3
23	10 24 41.0	68 42 35	46	1550	—	0.11	1
24	10 24 42.0	68 40 18	46	325	—	0.17	1
25	10 24 44.0	68 35 57	25	775	13	0.05	3
26	10 24 48.5	68 37 08	33	1240	—	0.07	1
27	10 24 49.0	68 41 09	48	744	—	0.17	1
28	10 24 49.5	68 39 43	53	279	6	0.17	3
29	10 24 52.5	68 43 22	56	356	10	0.09	3
30	10 24 56.0	68 39 59	53	1085	—	0.17	1
31	10 24 57.0	68 43 19	51	186	—	0.12	1
32	10 25 00.5	68 41 43	61	604	24	0.17	3
33	10 25 01.5	68 42 49	66	744	23	0.15	2
34	10 25 02.5	68 42 26	48	232	9	0.16	3
35	10 25 03.0	68 43 33	64	1023×496,74°	25	0.13	2
36	10 25 05.0	68 44 55	82	418	10	0.07	3
37	10 25 05.5	68 44 19	77	511	—	0.07	1
38	10 25 06.5	68 40 28	69	310	—	0.15	1
39	10 25 08.0	68 41 39	64	697	—	0.17	1
40	10 25 11.5	68 44 01	77	232	10	0.12	3
41	10 25 13.0	68 38 45	59	837	—	0.06	1
42	10 25 14.5	68 43 57	77	279	13	0.13	2
43	10 25 15.0	68 37 57	56	697×418,80°	—	0.05	2
44	10 25 16.5	68 37 53	56	186	8	0.05	3
45	10 25 18.5	68 41 13	87	418	7	0.16	3
46	10 25 21.8	68 41 57	87	155	4	0.16	3
47	10 25 28.5	68 41 13	71	155	8	0.05	3
48	10 25 34.0	68 42 54	92	403×186,80°	8	0.15	2

Table 4: Derived properties of the H I holes

#	d (pc)	v_{exp} (km s ⁻¹)	age t (10 ⁶ yr)	mass M_{HI} (10 ⁴ M _⊙)	energy E_{E} (10 ⁵⁰ ergs)
1	310	8	19.6	4.9	6.8
2	310	5	31.3	4.9	3.5
3	976.5	17	29.0	178.4	817.9
4	1178	14	42.5	104.4	326.9
5	775	9	43.5	59.5	103.7
6	325.5	8	20.6	1.8	2.1
7	1162.5	—	—	321.1	—
8	372	15	12.2	5.2	16.7
9	418.5	7	30.2	15.0	18.1
10	418.5	—	—	14.0	—
11	697.5	8	44.1	17.3	22.7
12	325.5	—	—	7.0	—
13	1922	—	—	1451.2	—
14	325.5	—	—	7.5	—
15	697.5	6	58.7	69.4	71.6
16	325.5	8	20.6	7.0	9.9
17	465	13	18.1	10.3	27.4
18	434	11	19.9	17.8	40.7
19	232.5	8	14.7	1.1	1.4
20	837	—	—	119.9	—
21	604.5	18	17.0	48.0	228.3
22	744	13	28.9	89.4	276.7
23	1550	—	—	523.3	—
24	325.5	—	—	7.5	—
25	775	13	30.1	29.7	79.8
26	1240	—	—	170.5	—
27	744	—	—	89.4	—
28	279	6	23.5	4.7	4.4
29	356.5	10	18.0	5.2	9.5
30	1085	—	—	277.4	—
31	186	—	—	1.0	—
32	604.5	24	12.7	48.0	341.5
33	744	23	16.3	78.9	534.6
34	232.5	9	13.1	2.6	4.1
35	713	25	14.3	60.2	420
36	418.5	10	21.1	6.6	11.8
37	511.5	—	—	12.0	—
38	310	—	—	5.7	—
39	697.5	—	—	73.7	—
40	232.5	10	11.7	1.9	3.4
41	837	—	—	44.9	—
42	279	13	10.8	3.6	9.6
43	542.5	—	—	10.2	—
44	186	8	11.7	0.4	0.5
45	418.5	7	30.2	15.0	18.1
46	155	4	19.6	0.8	0.4
47	155	8	9.8	0.2	0.3
48	279	8	17.6	4.2	5.7

Table 5: Cross identification of our grouping of H II regions with the ones defined by MH94

Number	$\alpha_{1950.0}$	$\delta_{1950.0}$	cross identifications Miller & Hodge 1994 (MH94)
1'	10 25 20.2	68 41 40	246,251
2'	10 25 10.1	68 40 50	181,190,193,197,203,206, 207,218,221,226,227,230
3'	10 25 10.2	68 40 32	208
4'	10 25 10.4	68 40 19	195,213,214,227
5'	10 25 05.7	68 41 15	184
6'	10 25 04.5	68 39 39	182,174
7'	10 24 59.5	68 41 51	136,143,155,158,165
8'	10 24 54.1	68 40 53	125,126,131
9'	10 24 48.9	68 40 31	117
10'	10 24 49.9	68 40 18	120
11'	10 24 49.5	68 39 43	115,121
12'	10 24 48.7	68 39 27	113,116
13'	10 24 41.3	68 38 58	99,100,106
14'	10 24 44.5	68 40 07	111
15'	10 24 42.0	68 40 37	105
16'	10 24 40.1	68 40 05	94,96,102,104
17'	10 24 33.1	68 39 33	76,80,83,84
18'	10 24 20.4	68 37 55	57,58,60-63,65,67
19'	10 24 13.9	68 37 44	50
20'	10 24 09.3	68 37 24	46
21'	10 23 57.0	68 38 37	17,19,20
22'	10 24 02.7	68 39 12	30,33
23'	10 24 02.5	68 39 23	31
24'	10 24 00.1	68 39 17	25
25'	10 24 04.5	68 39 58	24,28,32,34,36,38-41,43-46
26'	10 23 59.7	68 39 58	24
27'	10 24 49.1	68 43 30	118
28'	10 24 58.4	68 42 29	144
29'	10 25 05.7	68 42 35	186
30'	10 25 03.9	68 43 04	178
31'	10 24 59.4	68 43 23	148
32'	10 24 56.1	68 43 24	133,134,137,138
33'	10 24 58.0	68 43 52	140
34'	10 25 03.0	68 43 51	167,169,179,180
35'	10 25 07.2	68 43 26	192,196
36'	10 25 08.4	68 43 49	194,198,202
37'	10 25 12.0	68 44 06	210,215,224,225
38'	10 25 14.8	68 43 51	235
39'	10 25 17.9	68 43 51	237,238,240,242,244,245
40'	10 25 15.7	68 43 20	233,237,240

Table 6: Derived properties of grouped H II regions

Number	$F(\text{H}\alpha)$ [$10^{-15} \text{ erg cm}^{-2} \text{ s}^{-1}$]	ρ_{HII} [cm^{-3}]	$r_{\text{HI,eff}}$ [pc]	M_{HI} [$10^5 M_{\odot}$]	Δv km s^{-1}	ρ_{HI} [cm^{-3}]	σ_{HI} [10^{20} cm^{-2}]	remarks
1'	30.8	0.7	201	14	16.3	1.68	13.9	HI-clump
2'	434.2	1.8	225	24	26.1	2.05	18.9	HI-clump
3'	32.9	0.85	117	5.8	26.3	3.52	16.9	new tiny hole ($D < 60 \text{ pc}$), rim hole 38
4'	41.9	0.8	148	10	19.7	3.00	18.3	diffuse HI
5'	13.7	0.5	152	7.4	22.4	2.05	12.8	on rim of hole 39, diffuse HI
6'	34.5	0.8	245	21	18.3	1.39	14.0	HI-clump
7'	152.4	0.6	270	17	22.4	0.84	9.3	H α fills hole 32!
8'	55.2	0.7	196	14	23.7	1.81	14.6	on rim of hole 27
9'	5.2	0.8	111	4.8	26.3	3.41	15.6	–
10'	5.2	0.6	107	4.7	21.7	3.73	16.4	on rim of hole 30
11'	40.8	0.6	100	3.9	28.9	5.23	15.6	H α -em. on rim of hole 28
12'	9.9	0.65	103	4.4	25.0	3.92	16.6	new tiny hole ($D < 60 \text{ pc}$)
13'	44.7	0.6	295	28	23.7	1.06	12.9	on rim of hole 22, diffuse HI
14'	8.9	0.9	88	2.7	26.3	3.85	13.9	on rim of hole 24, diffuse HI
15'	9.9	0.6	155	7.9	20.0	2.06	13.1	on rim of hole 24. diffuse HI
16'	102.2	0.9	163	10	34.2	2.25	15.0	on rim of holes 21,22,24, high Δv !
17'	86.8	1.4	142	8.5	25.0	2.89	16.9	on rim of holes 21,22
18'	83.8	0.8	303	41	25.0	1.43	18.7	a lot of HI in region
19'	139.0	1.1	130	8.1	20.4	3.58	19.1	a lot of HI in region
20'	13.6	0.7	111	4.3	22.4	3.00	14.0	a lot of HI in region
21'	75.4	0.85	233	17	18.4	1.30	12.5	new tiny hole ($D < 60 \text{ pc}$)
22'	27.5	1.0	77	2.4	19.6	5.11	16.2	HI-clump between H α -em.
23'	13.8	0.8	95	3.2	19.7	2.37	14.2	diffuse HI
24'	9.6	0.7	93	2.8	21.0	2.43	12.9	diffuse HI
25'	213.2	0.8	270	23	31.6	1.14	12.6	H α same as rim of hole 8!
26'	16.2	0.8	73	2.4	31.3	6.00	18.0	on rim of holes 8,13, high Δv
27'	35.9	0.9	124	6.2	25.0	3.16	16.1	on rim of holes 23,29
28'	7.8	0.6	89	2.7	21.0	3.73	13.6	new tiny hole ($D < 60 \text{ pc}$), rim hole 33
29'	109.0	0.6	93	4.1	26.3	4.96	18.9	diffuse HI, on rim of hole 33
30'	55.0	0.8	136	7.1	23.7	2.75	15.3	new tiny hole ($D < 60 \text{ pc}$)
31'	120.9	1.4	45	3.3	34.2	3.52	6.5	hardly any HI, high Δv
32'	204.3	1.5	175	14	33.0	2.54	18.3	on rim of holes 31,35, high Δv
33'	78.8	0.85	93	2.5	32.0	3.02	11.5	little HI, high Δv
34'	1004.7	2.0	199	8.9	25.0	1.03	9.0	little HI, on rim of holes 35,37
35'	444.9	2.0	154	11	23.5	2.93	18.5	a lot HI, on rim of hole 35
36'	487.2	1.9	183	17	28.7	2.70	20.3	a lot HI, rim hole 35, high Δv
37'	140.4	1.3	172	13	35.3	2.48	17.5	little HI, rim hole 40, high Δv ,
38'	13.3	0.8	98	5.0	23.7	5.17	20.8	on rim of hole 42
39'	144.8	1.2	207	28	22.4	3.07	26.1	a lot of HI in region
40'	15.4	0.85	182	15	25.6	2.42	18.1	a lot of HI in region

Table 7: Some properties of high mass stars (based on Devereux *et al.* 1997)

Type	Mass ($M_{\odot}$)	L_{bol} ($L_{\odot}$)	N_{Lyc}	$L(\text{H}\alpha)$	$L_{\text{bol}}/L(\text{H}\alpha)$
O5	40	6.8×10^5	4×10^{49}	1.4×10^4	49
O7	25	1.0×10^5	4×10^{48}	1.4×10^3	71
O9	20	4.6×10^4	1.3×10^{48}	4.6×10^2	100
B0	16	2.5×10^4	2.5×10^{47}	89	282
B1	10	5.2×10^3	1.9×10^{45}	0.67	7.8×10^3
B2	8	2.9×10^3	4.5×10^{44}	0.16	1.8×10^4
B5	6	7.9×10^2	4.8×10^{43}	0.02	4.0×10^4

Table 8: Summary of H I hole statistics in four nearby galaxies

Property	M 31 ^a	M 33 ^b	IC 2574 ^c	Ho II ^d
Linear resolution (pc)	100	55	95	65
Velocity resolution (km s^{-1})	8.2	8.2	2.6	2.6
Average surface density (10^{20} cm^{-2})	5	9	4	10
Average volume density (cm^{-3})	0.6	0.45	0.15	0.2
Average velocity dispersion (km s^{-1})	8	8	7	7
Derived scaleheight (pc)	120	100	350	625
Number of holes	141	148	48	51
Sensitivity per channel (10^{20} cm^{-2})	0.3	1.0	0.5	2.4

^aBrinks & Bajaja 1986

^bDeul & den Hartog 1990

^cThis paper

^dPuche *et al.* 1992

This figure "walter.fig1n.jpeg" is available in "jpeg" format from:

<http://arXiv.org/ps/astro-ph/9904002v1>

This figure "walter.fig2n.jpeg" is available in "jpeg" format from:

<http://arXiv.org/ps/astro-ph/9904002v1>

This figure "walter.fig3an.jpeg" is available in "jpeg" format from:

<http://arXiv.org/ps/astro-ph/9904002v1>

This figure "walter.fig3bn.jpeg" is available in "jpeg" format from:

<http://arXiv.org/ps/astro-ph/9904002v1>

This figure "walter.fig3cn.jpeg" is available in "jpeg" format from:

<http://arXiv.org/ps/astro-ph/9904002v1>

This figure "walter.fig4n.jpeg" is available in "jpeg" format from:

<http://arXiv.org/ps/astro-ph/9904002v1>

This figure "walter.fig5n.jpeg" is available in "jpeg" format from:

<http://arXiv.org/ps/astro-ph/9904002v1>

This figure "walter.fig6n.jpeg" is available in "jpeg" format from:

<http://arXiv.org/ps/astro-ph/9904002v1>

This figure "walter.fig7n.jpeg" is available in "jpeg" format from:

<http://arXiv.org/ps/astro-ph/9904002v1>

This figure "walter.fig8n.jpeg" is available in "jpeg" format from:

<http://arXiv.org/ps/astro-ph/9904002v1>

This figure "walter.fig9n.jpeg" is available in "jpeg" format from:

<http://arXiv.org/ps/astro-ph/9904002v1>

This figure "walter.fig10n.jpeg" is available in "jpeg" format from:

<http://arXiv.org/ps/astro-ph/9904002v1>

This figure "walter.fig11n.jpeg" is available in "jpeg" format from:

<http://arXiv.org/ps/astro-ph/9904002v1>

This figure "walter.fig12n.jpeg" is available in "jpeg" format from:

<http://arXiv.org/ps/astro-ph/9904002v1>

This figure "walter.fig13n.jpeg" is available in "jpeg" format from:

<http://arXiv.org/ps/astro-ph/9904002v1>

This figure "walter.fig14n.jpeg" is available in "jpeg" format from:

<http://arXiv.org/ps/astro-ph/9904002v1>

This figure "walter.fig15n.jpeg" is available in "jpeg" format from:

<http://arXiv.org/ps/astro-ph/9904002v1>

This figure "walter.fig16n.jpeg" is available in "jpeg" format from:

<http://arXiv.org/ps/astro-ph/9904002v1>

This figure "walter.fig17n.jpeg" is available in "jpeg" format from:

<http://arXiv.org/ps/astro-ph/9904002v1>

This figure "walter.fig18n.jpeg" is available in "jpeg" format from:

<http://arXiv.org/ps/astro-ph/9904002v1>

This figure "walter.fig19n.jpeg" is available in "jpeg" format from:

<http://arXiv.org/ps/astro-ph/9904002v1>

This figure "walter.fig20n.jpeg" is available in "jpeg" format from:

<http://arXiv.org/ps/astro-ph/9904002v1>